arXiv:hep-ph/0607178v2 20 Oct 2006

Calculation of BR($\overline{B}^0 \to \Lambda_c^+ + \bar{p}$) in the PQCD Approach

Xiao-Gang He[1,2], Tong Li[1], Xue-Qian Li[1], and Yu-Ming Wang[1]

[1]Department of Physics, Nankai University, Tianjin

[2]Department of Physics and Center for Theoretical Sciences, National Taiwan University, Taipei

Abstract:

We calculate the branching ratio of $\overline{B}^0 \to \Lambda_c^+ \overline{p}$ in the PQCD approach. Most previous model calculations obtained branching ratios significantly larger than experimental data. We find that the predicted branching ratio for BR($\overline{B}^0 \to \Lambda_c^+ \overline{p}$) in the PQCD approach can vary over a range of $(2.3 \sim 5.1) \times 10^{-5}$ with the largest uncertainty coming from the parameters in the wave function of Λ_c. With the favored values for the parameters in the Λ_c^+ wave function, $\beta = 1$ GeV and $m_q = 0.3$ GeV, the branching ratio is about 2.3×10^{-5} which is satisfactorily consistent with the value measured by experiments.

PACS numbers: 13.30.Ce, 12.38.Bx, 14.20.Mr

I. INTRODUCTION

B-physics stands as an ideal laboratory for studying hadron structure and fundamental mechanisms which govern the reactions for a heavy quark and the related hadronization processes. It also offers a window to search for new physics. The mesonic decays of B-meson have already been carefully studied by many authors because such processes are theoretically simpler and have a rich data accumulation. Nevertheless much data on the baryonic decays of B-mesons have also been collected. In 1987, $B \to p\overline{p}\pi^{\pm}$ and $B \to p\overline{p}\pi^{+}\pi^{-}$ were firstly observed by ARGUS[1] and since then, various baryonic B baryonic-decay modes were measured at CLEO, Belle and BaBar[2, 3, 4, 5, 6]. All the information offers us an opportunity to more seriously investigate the processes where baryons are involved. However, the case for baryonic decays is much more complicated than the mesonic one, especially in the perturbative QCD (PQCD) approach[7] where there is not any real spectator constituent, namely all the constituents must be connected by exchanged gluons. One also needs to have detailed information for the wave functions of baryons which contain three valence quarks. All these make the problem much more complicated and cause theoretical uncertainties.

In this work we study the process $\overline{B}^0 \to \Lambda_c^+\overline{p}$. This process has been measured [3, 4, 5] with a branching ratio of [6] $(2.2{\pm}0.8){\times}10^{-5}$. There have been some theoretical evaluations for this branching ratio using various phenomenological models, such as the constituent quark model, the pole model, the QCD sum rule, the diquark model and others[8, 9, 10, 11, 12]. All the theoretical predictions made in these models are substantially larger than the data. Later Cheng and Yang[13] considered the pole structure and applied the bag model approach for calculating the hadronic matrix element. Their result is close to the experimental data. The related theoretical estimations on the branching ratio and the experimental data are listed in Table I.

Since the success of the PQCD approach in studying mesonic decays of B-meson is remarkable, it is natural to extend this approach to evaluate the decay rates of baryonic decays of B-mesons. In $\overline{B}^0 \to \Lambda_c^+\overline{p}$, the two final baryons are much lighter than B-meson, they possess relatively large linear momentum $\mathbf{p} = \lambda^{1/2}(M_B^2, M_{\Lambda_c}^2, M_p^2)/2M_B$ in the rest frame of B-meson, thus the momentum match of the quarks in the final baryons requires that the exchanged gluons must be "hard", or say, have large k^2. Here k indicates the generic momentum carried by a gluon. Moreover, unlike in some cases where a pair of quark-antiquark is created from vacuum, the pair of quark-antiquark is created by a hard gluon, therefore one expects that the whole process is calculable in the framework of PQCD[14].

In the PQCD approach, there are unknown parameters in the wave functions of the B-meson and the two baryons in the final state. However, since these parameters are universal, they can be fixed by fitting data of other relevant decays. This treatment of the unknown parameters reduces the model dependence of the theoretical calculations. Therefore one has good reasons to believe that the result obtained in PQCD would be credible. Indeed, our numerical result is closer to the experimental data than those by other models.

TABLE I: Branching ratio of $\overline{B}^0 \to \Lambda_c^+\overline{p}$ for different models and experiments (last two numbers), respectively. There is also a preliminary Babar result[4] $(2.15 \pm 0.36 \pm 0.13 \pm 0.56) \times 10^{-5}$.

$\overline{B}^0 \to \Lambda_c^+\overline{p}$	[9]	[10]	[11]	[12]	[13]	[3]	[5]
BR	4×10^{-4}	8.5×10^{-4}	1.1×10^{-3}	$(1.7\sim1.9)\times10^{-3}$	1.1×10^{-5}	$(2.19^{+0.56}_{-0.49}\pm0.32\pm0.57)\times10^{-5}$	$<9\times10^{-5}$

The effective Hamiltonian responsible at the quark level for $\overline{B}^0 \to \Lambda_c^+\overline{p}$ is given by[15]:

$$H_{eff} \;=\; \frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^{*}\left[C_1(\overline{c}b)_{V-A}(\overline{d}u)_{V-A}+C_2(\overline{c}u)_{V-A}(\overline{d}b)_{V-A}\right]+h.c., \tag{1}$$

where $(\bar{a}b)_{V-A} = \bar{a}\gamma^{\mu}(1-\gamma_5)b$, and $C_1^{LO}(m_b) = 1.12, C_2^{LO}(m_b) = -0.27$[15]. In our calculation the μ dependence of $C_{1,2}$ at the leading order remains.

At the hadron level, the decay amplitude for $\overline{B}^0 \to \Lambda_c^+\overline{p}$ is obtained by sandwiching the effective Hamiltonian between the initial and final hadron states,

$$\begin{aligned} M(\overline{B}^0 \to \Lambda_c^+\overline{p}) &= \langle \Lambda_c^+\overline{p}|H_{eff}|\overline{B}^0\rangle \\ &= \frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^{*}\overline{\Lambda_c^+}\left(A+B\gamma_5\right)p, \end{aligned} \tag{2}$$

where A and B are two form factors which we will calculate in the PQCD approach.

The partial decay width is then given by

$$\Gamma(\overline{B}^0 \to \Lambda_c^+ \overline{P}) = \frac{G_F^2 |V_{cb}|^2 |V_{ud}|^2}{16 m_B^3 \pi} \sqrt{(m_B^2 - (m_{\Lambda_c} + m_P)^2)(m_B^2 - (m_{\Lambda_c} - m_P)^2)} \\ \left[|A|^2 (m_B^2 - (m_{\Lambda_c} + m_P)^2) + |B|^2 (m_B^2 - (m_{\Lambda_c} - m_P)^2)\right] . \tag{3}$$

In the following sections we present details for the calculations. In Section II, we describe the approach for calculations. In Section III, we present our numerical results along with all necessary input parameters. The last section is devoted to our discussions and conclusions. Detailed calculations for each diagram are given in the Appendices.

II. PQCD CALCULATION OF THE HADRONIC MATRIX ELEMENTS

The effective Hamiltonian at quark level has been studied by many authors and the PQCD approach has also extensively been discussed in literature. In this section we discuss how to calculate the hadronic matrix elements defined in the previous section in terms of the PQCD method[7, 16, 17, 18]. As usual, we define, in the rest frame of $\overline{B^0}$, q, p and p' to be the four-momenta of $\overline{B^0}$, Λ_c^+ and antiproton, q is the b quark momentum, $q_i (i = 1, 2)$, $k_i (i = 1, 2, 3)$ and k_i' to be the momenta of the valance light quarks (anti-quark) inside $\overline{B^0}$, Λ_c^+, and $\overline{p}$ respectively. We parameterize the corresponding light cone momenta with the masses of all light quarks and antiproton being neglected as

$$\begin{aligned}
&q = (q^+, q^-, \mathbf{0}_T) = \frac{m_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), \ \ p = (p^+, p^-, \mathbf{0}_T) = (q^+, r^2 q^-, \mathbf{0}_T) = \frac{m_B}{\sqrt{2}}(1, r^2, \mathbf{0}_T) \\
&p' = (0, p'^-, \mathbf{0}_T) = \frac{m_B}{\sqrt{2}}(0, 1 - r^2, \mathbf{0}_T) \\
&q_1 = (y q^+, q^-, \mathbf{q}_T), \ \ q_2 = ((1 - y) q^+, 0, -\mathbf{q}_T) \\
&k_1 = (x_1 p^+, p^-, \mathbf{k}_{1T}), \ \ k_2 = (x_2 p^+, 0, \mathbf{k}_{2T}), \ \ k_3 = (x_3 p^+, 0, \mathbf{k}_{3T}) \\
&k_1' = (0, x_1' p'^-, \mathbf{k}_{1T}'), \ \ k_2' = (0, x_2' p'^-, \mathbf{k}_{2T}'), \ \ k_3' = (0, x_3' p'^-, \mathbf{k}_{3T}')
\end{aligned} \tag{4}$$

where $r = m_{\Lambda_c}/m_B$, and y, x_i, x_i' are the fractions of the longitudinal momenta of the valence quarks with $x_1 + x_2 + x_3 = 1$ and $x_1' + x_2' + x_3' = 1$. $\mathbf{q}_T$, $\mathbf{k}_{iT}$ and $\mathbf{k}_{iT}'$ are the transverse momenta of the valence quarks inside $\overline{B^0}$, Λ_c^+ and antiproton, respectively.

Note that because $\overline{B^0}$ is in its rest frame, even though the momenta of the valance quarks inside the final states $(\Lambda_c^+, \overline{p})$ are fixed, the four-momenta of the valance quarks inside $\overline{B^0}$ (q_1, q_2) are not uniquely determined. One can also choose to have:

$$q_1 = (q^+, y q^-, \mathbf{q}_T), \ \ q_2 = (0, (1 - y) q^-, -\mathbf{q}_T). \tag{5}$$

Obviously, the first case (q_i^1) corresponds to that the momenta of $\overline{d}$-quarks in initial state $\overline{B^0}$ and in final state $\overline{p}$ are opposite to each other, and the second one (q_i^2) is in the same direction as the antiproton momentum (i.e. they are parallel.). One has to convolute the initial light-cone wave functions with the final light-cone wave function to make the correct choice.

In the PQCD picture, hadrons are made of valence quarks whose momenta-distributions are described by appropriate wave functions. The wave function of Λ_c is usually defined through the correlator[19],

$$\begin{aligned}
(Y_{\Lambda_c})_{\alpha\beta\gamma}(k_i, \nu) &= \frac{1}{2\sqrt{2} N_c} \int \prod_{l=2}^{3} \frac{dw_l^+ d\mathbf{w_l}}{(2\pi)^3} e^{i k_l w_l} \varepsilon^{abc} \langle 0 | T[c_\alpha^a(0) u_\beta^b(w_2) d_\gamma^c(w_3)] | \Lambda_c(p) \rangle \\
&= \frac{f_{\Lambda_c}}{8\sqrt{2} N_c} [(\not{p} + m_{\Lambda_c}) \gamma_5 C]_{\beta\gamma} [\Lambda_c(p)]_\alpha \Psi(k_i, \nu),
\end{aligned} \tag{6}$$

where f_{Λ_c} is a normalization constant, $\Lambda_c(p)$ is the Λ_c spinor, and $\Psi(k_i, \nu)$ is the wave function. ν is the hard sub-amplitude energy scale for the process which is within the range of $\Lambda_{QCD} < \nu < m_b$ and will be further discussed below. Similarly, the leading-twist wave function of proton is defined by[16, 20]:

$$
\begin{aligned}
&(Y_P)_{\alpha\beta\gamma}(k'_i,\nu) = \frac{1}{2\sqrt{2}N_c}\int\prod_{l=1}^{2}\frac{dw_l^- d\mathbf{w_l}}{(2\pi)^3}e^{ik'_l w_l}\varepsilon^{abc}\langle 0|T[u^a_\alpha(w_1)u^b_\beta(w_2)d^c_\gamma(0)]|P(p')\rangle \\
&= \frac{f_P(\nu)}{8\sqrt{2}N_c}\{(\not{p}'C)_{\beta\gamma}[\gamma_5 P(p')]_\alpha\Phi^V(k'_i,\nu) + (\not{p}'\gamma_5 C)_{\beta\gamma}[P(p')]_\alpha\Phi^A(k'_i,\nu) \\
&-(\sigma_{\mu\nu}p'^\nu C)_{\beta\gamma}[\gamma^\mu\gamma_5 P(p')]_\alpha\Phi^T(k'_i,\nu)\},
\end{aligned}
\tag{7}
$$

where f_P is the normalization constant, and $P(p')$ is the proton spinor.

$$
\begin{aligned}
\Phi^V(k'_1,k'_2,k'_3,\nu) &= \frac{1}{2}\left[\Phi(k'_2,k'_1,k'_3,\nu)+\Phi(k'_1,k'_2,k'_3,\nu)\right]\ , \\
\Phi^A(k'_1,k'_2,k'_3,\nu) &= \frac{1}{2}\left[\Phi(k'_2,k'_1,k'_3,\nu)-\Phi(k'_1,k'_2,k'_3,\nu)\right]\ , \\
\Phi^T(k'_1,k'_2,k'_3,\nu) &= \frac{1}{2}\left[\Phi(k'_1,k'_3,k'_2,\nu)+\Phi(k'_2,k'_3,k'_1,\nu)\right]\ .
\end{aligned}
\tag{8}
$$

The B meson wave function is expressed as[21]

$$
\begin{aligned}
(Y_B)_{\alpha\beta}(q_i,\nu) &= \frac{1}{N_c}\int\frac{d^4w}{(2\pi)^4}e^{iq_2\cdot w}\langle 0|T[\overline{d}_\alpha(w)b_\beta(0)]|\overline{B^0}(q)\rangle \\
&= \frac{i}{\sqrt{2N_c}}\left[(\not{q}+m_B)\gamma_5\left(\frac{\not{n}_+}{\sqrt{2}}\phi_B^+(q_2,\nu)+\frac{\not{n}_-}{\sqrt{2}}\phi_B^-(q_2,\nu)\right)\right]_{\beta\alpha}.
\end{aligned}
\tag{9}
$$

where $\not{n}_+$ term corresponds to the first choice of momenta of the valance quark $\overline{d}$ in $\overline{B^0}$ and $\not{n}_-$ term corresponds to the second one with the relation:

$$
\frac{\not{n}_\pm}{\sqrt{2}} = \frac{\not{q}_2^{1,2}}{(1-y)m_B}.
\tag{10}
$$

When convoluting the hadron wave functions with the hard amplitude, one should use corresponding light cone momenta $q_i^{1,2}$ consistently for the above two terms.

There is also a simple relation between wave functions ϕ_B^+, ϕ_B^- and $\phi_B, \overline{\phi}_B$ used in the usual PQCD calculation:

$$
\phi_B^+ = \phi_B + \overline{\phi}_B, \qquad \phi_B^- = \phi_B - \overline{\phi}_B
\tag{11}
$$

It has been argued that the distribution amplitude $\overline{\phi}_B$ is negligible compared to ϕ_B[21], therefore $\phi_B^+ = \phi_B^- \simeq \phi_B$. The specific form of ϕ_B will be given later when carrying our numerical calculations.

Including the Sudakov factor with infrared cut-offs $\omega(\omega',\omega_q)$, and running the wave function from ν down to $\omega(\omega',\omega_q)$, the wave functions are obtained as [16]:

$$
\begin{aligned}
\Psi(x_i,b_i,p,\nu) &= \exp\left[-\sum_{l=2}^{3}s(\omega,x_l p^+) - 3\int_\omega^\nu\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))\right]\Psi(x_i) \\
\Phi(x'_i,b'_i,p',\nu) &= \exp\left[-\sum_{l=1}^{3}s(\omega',x'_l p^-) - 3\int_{\omega'}^\nu\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))\right]\Phi(x'_i,\omega'), \\
\Phi_B(y,b_q,q,\nu) &= \exp\left[-s(\omega_q,q_2^+) - 2\int_{\omega_q}^\nu\frac{d\bar{\mu}}{\bar{\mu}}\gamma(\alpha_s(\bar{\mu}))\right]\Phi_B(1-y)
\end{aligned}
\tag{12}
$$

where $\omega = min(1/\tilde{b}_1, 1/\tilde{b}_2, 1/\tilde{b}_3)$, $\omega' = min(1/\tilde{b}'_1, 1/\tilde{b}'_2, 1/\tilde{b}'_3)$ and $\omega_q = 1/b_q$. $\tilde{b}_1^{(')} = |\mathbf{b}_2^{(')} - \mathbf{b}_3^{(')}|$, $\tilde{b}_2^{(')} = |\mathbf{b}_1^{(')} - \mathbf{b}_3^{(')}|$, $\tilde{b}_3^{(')} = |\mathbf{b}_1^{(')} - \mathbf{b}_2^{(')}|$ and $b_q = |\mathbf{b}_q|$. Here $\mathbf{b}$, $\mathbf{b}'$ and $\mathbf{b}_q$ are the conjugate variables to $\mathbf{k}_T$, $\mathbf{k}'_T$ and $\mathbf{q}_T$ whose definitions are collected in Appendix B.

The explicit expressions for the Sudakov factors are given in [16] with

$$
\begin{aligned}
&s(\omega, Q) = \int_{\omega}^{Q} \frac{dp}{p}\left[ln(\frac{Q}{p})A[\alpha_s(p)] + B[\alpha_s(p)]\right] \\
&A = C_F\frac{\alpha_s}{\pi} + \left[\frac{67}{9} - \frac{\pi^2}{3} - \frac{10}{27}n_f + \frac{8}{3}\beta_0 ln(\frac{e^{\gamma_E}}{2})\right](\frac{\alpha_s}{\pi})^2 \\
&B = \frac{2}{3}\frac{\alpha_s}{\pi} ln\left(\frac{e^{2\gamma_E - 1}}{2}\right) \\
&\gamma_q(\alpha_s(\mu)) = -\alpha_s(\mu)/\pi, \\
&\beta_0 = \frac{33 - 2n_f}{12},
\end{aligned}
\tag{13}
$$

where γ_E is the Euler constant, n_f is the flavor number and γ_q is the anomalous dimension. For B decays, the typical energy scale is above the charm mass, so we will take n_f to be 4 in our numerical computations.

Within the PQCD framework, the hadronic matrix element can be written as:

$$
\begin{aligned}
&M = \int[Dx]\int[Db](\overline{Y}_{\Lambda_c})_{\alpha'\beta'\gamma'}(x'_i, b'_i, p', \nu) \\
&H^{\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x'_i, y, b_i, b'_i, b_q, m_B, m_{\Lambda_c}, \nu)(Y_{\overline{p}})_{\alpha\beta\gamma}(x_i, b_i, p, \nu)(Y_B)_{\rho\rho'}(y, b_q, q, \nu),
\end{aligned}
\tag{14}
$$

where

$$
[Dx] = [dx][dx']dy, \;\; [dx] = dx_1 dx_2 dx_3 \delta(1 - \sum_{l=1}^{3} x_l), \;\; [dx'] = dx'_1 dx'_2 dx'_3 \delta(1 - \sum_{l=1}^{3} x'_l).
\tag{15}
$$

The measures of the transverse parts $[Db]$ are defined in Appendix A.

To obtain the hard scattering amplitude $H^{\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x, x', y, b, b', b_q, m_B, m_{\Lambda_c}, \nu)$, one can derive the amplitude $H^{i,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x'_i, y, \mathbf{k}_T, \mathbf{k}'_T, \mathbf{q}_T, m_B, m_{\Lambda_c})$ corresponding to the 'i'th diagram in Fig. 1 and its analytic expression is displayed in Appendix B. Then a Fourier transformation on $\mathbf{k}_T$ and $\mathbf{k}'_T$ is carried out to convert them into the $\mathbf{b}$ and $\mathbf{b}'$ space to obtain $\tilde{H}^{i,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x, x', y, b, b', b_q, m_B, m_{\Lambda_c})$. The concrete procedure for carrying out such transformation is described at the end of Appendix B.



FIG. 1: The lowest order diagrams for the $\overline{B}^0 \rightarrow \Lambda_c^+\overline{p}$ decay according to the first choice of momenta (q_i^1). The wavy line indicates W-exchange. The solid lines, curly lines denote the quarks and gluons, respectively. In the diagrams, only one gluon line is shown. The other gluon line connects the solid black blob and the cross indicated by a_i. There are 36 diagrams. We label each one by a_i.

We can catalog the diagrams into several types. The first type includes 36 diagrams, indicated by a_i in Fig. 1. If we adopt the first choice of momenta (q_i^1) of valance quark in $\overline{B^0}$ the total 36 diagrams contribute to the amplitude



FIG. 2: The lowest order diagrams for the $\overline{B}^0 \to \Lambda_c^+ \overline{p}$ decay according to the second choice of momenta (q_i^2). The wavy line indicates W-exchange. The solid lines, curly lines denote the quarks and gluons, respectively. There are 8 diagrams. We label each one by b_i.



FIG. 3: The lowest order diagrams for the $\overline{B}^0 \to \Lambda_c^+ \overline{p}$ decay including three-gluon vertices. The wavy line indicates W-exchange. The solid lines, curly lines denote the quarks and gluons, respectively. There are 4 diagrams. We label each one by c_i.



FIG. 4: The W-exchange diagrams for the $\overline{B}^0 \to \Lambda_c^+ \overline{P}$ decay.

in the PQCD frame, where the gluons can be "hard" (have large k^2). However, one can notice that, if we adopt the second choice (q_i^2), in some of the diagrams the gluon attached to the spectator quark is always soft, i.e. $k^2 \approx 0$. The contributions of such diagrams cannot be counted as PQCD contributions and are attributed into the wave functions of the initial or final hadrons. When calculating the PQCD contributions, these diagrams should not be included at all. In that case, there are only 8 diagrams, indicated by b_i in Fig. 2. remain where all the exchanged gluons are "hard". For another type of diagrams, 4 of them, indicated by c_i in Fig. 3, three-gluon vertices are involved. We find that the color factor for two, $c_{3,4}$, of them is zero. There are also the W-exchange diagrams (where W is exchanged between the initial b and $\bar{d}$), 36 of them with gluons connected to quark lines only and 4 of them with three-gluon vertices shown in Fig. 4. These diagrams are suppressed by the linear-momentum match and usually ignored[13]. Therefore, there are 36+8+2=46 diagrams shown in Fig. 1, 2 and 3, need to be evaluated to the leading order.

Multiplying all contributions of the diagrams in Fig.1, 2 and 3 with corresponding Wilson coefficients, one adds them up to obtain the full hard scattering amplitude

$$H^{\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x,x',y,b,b',b_q,m_B,m_{\Lambda_c})=\sum_i C(t)\tilde{H}^{i,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x,x',y,b,b',b_q,m_B,m_{\Lambda_c}).$$

Here we denote the hard scale as t which is taken to be the larger of the two variables $t_{1,2}$ associated with the virtual gluon momentum in Fig.1, i.e. $t=\max(t_1^i,t_2^i)$. The expressions for $t_{1,2}$ are listed in Appendix C.

Finally a RG running is applied to the hard scattering amplitude from the scale ν in the wavefunctions to t and we obtain

$$\begin{aligned}&H^{\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x,x',y,b,b',b_q,m_B,m_{\Lambda_c},\nu)=\\&\exp[-8\int_\nu^t\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))]\times H^{\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x,x',y,b,b',b_q,m_B,m_{\Lambda_b}).\end{aligned}\tag{16}$$

The form factors are obtained by properly grouping relevant terms according to the definition in eq.(2). Using eq.(14) we obtain a generic expression for the form factor corresponding to each diagram as

$$\begin{aligned}A^i(B^i)\ &=\ \sum_{j=V,A,T}\frac{1}{128N_c^2\sqrt{2N_c}}f_Pf_{\Lambda_c}\int[Dx]\int[Db]^iC^i(t^i)\\&\quad\Psi_{\Lambda_c}(x)\Phi_P^j(x')\phi_B(1-y)\exp[-S^i]H_F^{ij}\Omega^i,\end{aligned}\tag{17}$$

where

$$\begin{aligned}S^i\ &=\ \sum_{k=2}^3 s(\omega,x_kp^+)+\sum_{k=1}^3 s(\omega',x'_kp'^-)+s(\omega_q,(1-y)q^+)\\&\quad+3\int_\omega^{t^i}\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))+3\int_{\omega'}^{t^i}\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))+2\int_{\omega_q}^{t^i}\frac{d\overline{\mu}}{\overline{\mu}}\gamma_q(\alpha_s(\overline{\mu}))\end{aligned}\tag{18}$$

and A^i and B^i represent the form factors contributed by the i-th diagram. C^i is an appropriate combination of the Wilson coefficients. The superscript j labels different Lorentz structures V, A, and T of anti-proton according to eq.(7). The explicit expressions of Ω^i are presented in Appendix D and the functions H_F^{ij} are given in Appendix E. The total form factors are obtained by summing over the contributions of all the diagrams.

III. NUMERICAL RESULTS

We are now ready to evaluate the form factors numerically. In our calculations, we adopt the distribution amplitude Ψ proposed in Ref.[17, 19] for the Λ_c ,

$$\Psi(x_1,x_2,x_3)=Nx_1x_2x_3\exp[-\frac{m_{\Lambda_c}^2}{2\beta^2x_1}-\frac{m_q^2}{2\beta^2x_2}-\frac{m_q^2}{2\beta^2x_3}].\tag{19}$$

The normalization constant N is fixed by the condition:

$$\int[dx]\Psi(x_1,x_2,x_3)=1.\tag{20}$$

Since the decay constant f_{Λ_c} is not experimentally determined so far, one has to invoke a theoretical evaluation to fix it. It is assumed that there is a relation between f_{Λ_c} and f_{Λ_c} as $f_{\Lambda_c} = f_{\Lambda_b}\frac{m_{\Lambda_b}}{m_{\Lambda_c}}$[17]. In terms of the PQCD method f_{Λ_b} is determined by fitting $B(\Lambda_b \to \Lambda_c l\overline{\nu})$ whose central value is 5% measured by the DELPHI collaboration[22]. In our previous study of $\Lambda_b \to \Lambda\gamma$[23], we have carried out a new determination for f_{Λ_b}. When fitting the data we first truncate the double logarithmic Sudakov factor in such a way that we require the factor $\exp(-s)$ to be smaller than 1 following the prescription of Ref.[24]. It is noted that our numerical value of f_{Λ_b} is different from that obtained in Ref.[17], where $B(\Lambda_b \to \Lambda_c l\overline{\nu})$ was taken to be 2%, whereas we use the newly measured value instead. In concrete computations, we choose the cut-offs as $\omega = 1.14 min(1/\tilde{b}_1, 1/\tilde{b}_2, 1/\tilde{b}_3)$ and $\omega' = 1.14 min(1/\tilde{b}'_1, 1/\tilde{b}'_2, 1/\tilde{b}'_3)$. The phenomenological factor 1.14 is adopted according to Ref.[25] to make the whole picture more realistic. Moreover, the values of β and m_q in the heavy baryon wave function need to be fixed. In Ref.[16, 17, 18], $\beta = 1$ GeV and $m_q = 0.3$ GeV were used to estimate the decay rates of $\Lambda_b \to \Lambda_c l\bar{\nu}$, $\Lambda_b \to pl\bar{\nu}$ and $\Lambda_b \to \Lambda J/\psi$. It is commonly expected that β should not be too much smaller than 1 GeV if the form factors are dominated by the perturbative contributions. To see how different values of β and m_q affect the results, we will let both β and m_q vary within ranges of $0.6 \sim 1.2$ GeV and $0.2 \sim 0.3$ GeV respectively. Using the fitted value of f_{Λ_b} obtained in Ref.[23], and the relation: $f_{\Lambda_c} = f_{\Lambda_b}\frac{m_{\Lambda_b}}{m_{\Lambda_c}}$, we listed f_{Λ_c} in Table II for different parameter sets.

TABLE II: Decay constant f_{Λ_c}(GeV) for different choices of β and m_q with Sudkov truncation described in the text.

f_{Λ_c}	$\beta = 0.6$GeV	$\beta = 0.8$GeV	$\beta = 1$GeV	$\beta = 1.2$GeV
$m_q = 0.2$GeV	1.70×10^{-3}	2.50×10^{-3}	3.51×10^{-3}	4.71×10^{-3}
$m_q = 0.3$GeV	3.12×10^{-3}	4.07×10^{-3}	5.22×10^{-3}	6.50×10^{-3}

The proton distribution amplitudes have been studied using the KS model. In this work, we adopt the model proposed in Ref.[20],

$$\begin{aligned}
\Phi(x_i, \omega) &= \phi_{as}(x_1, x_2, x_3) \sum_{j=0}^{5} N_j \left[\frac{\alpha_s(\omega)}{\alpha_s(\mu_0)}\right]^{b_j/(4\beta_0)} a_j A_j(x_i), \\
\phi_{as}(x_1, x_2, x_3) &= 120 x_1 x_2 x_3. \qquad (21)
\end{aligned}$$

with $\mu_0 \approx 1$ GeV, the constants N_j, a_j, b_j and the Appel polynomials A_j are listed in Table III[16].

TABLE III: The constants and polynomials in eq.(21).

j	a_j	N_j	b_j	$A_j(x_i)$
0	1.00	1	0	1
1	0.310	21/2	20/9	$x_1 - x_3$
2	-0.370	7/2	24/9	$2 - 3(x_1 + x_3)$
3	0.630	63/10	32/9	$2 - 7(x_1 + x_3) + 8(x_1^2 + x_3^2) + 4x_1x_3$
4	0.00333	567/2	40/9	$x_1 - x_3 - \frac{4}{3}(x_1^2 - x_3^2)$
5	0.0632	81/5	42/9	$2 - 7(x_1 + x_3) + \frac{14}{3}(x_1^2 + x_3^2) + 14x_1x_3$

The constant f_P is fixed to be[20]

$$f_P(\omega) = f_P(\mu_0) \left[\frac{\alpha_s(\omega)}{\alpha_s(\mu_0)}\right]^{1/(6\beta_0)}, \qquad (22)$$

with $f_P(\mu_0) = (5.2 \pm 0.3) \times 10^{-3}$ GeV^2.

The B meson distribution amplitude is given as [21]:

$$\phi_B(x, b) = N_B x^2 (1 - x)^2 exp\left[-\frac{m_B^2 x^2}{2\omega_b^2} - \frac{1}{2}(\omega_b b)^2\right] \qquad (23)$$

where $\omega_b = 0.4$ GeV and $N_B = 91.74$ GeV for $f_B = 0.19$ GeV.

Finally, when obtaining the branching ratio of $\overline{B}^0 \to \Lambda_c^+ \overline{p}$, for definitiveness we fix rest of the parameters as following. The parameter Λ_{QCD} which enters in the strong coupling constant and various Wilson coefficients is set to be $\Lambda_{QCD} = 0.2$ GeV. For the CKM mixing parameters, we take their central values[6]: $s_{12} = 0.2243$, $s_{23} = 0.0413$, $s_{13} = 0.0037$ and $\delta_{13} = 1.05$. We take two typical values for m_q as 0.2 and 0.3 GeV and let β vary within a reasonable range to carry out the numerical computations.

The theoretical values of the branching ratio for different β and m_q are listed in Table IV. Our numerical results indicate that BR($\overline{B}^0 \to \Lambda_c^+ \overline{p}$) = $(2.3 \sim 5.1) \times 10^{-5}$. We see that the results are consistent with experimental data. With the values $\beta = 1$ GeV and $m_q = 0.3$ GeV used in Ref.[16, 17, 18], BR($\overline{B}^0 \to \Lambda_c^+ \overline{p}$) = 2.3×10^{-5}.

TABLE IV: Branching ratio($\times 10^{-5}$) for different choices of β and m_q with Sudakov truncation described in the text.

BR	$\beta = 0.6$GeV	$\beta = 0.8$GeV	$\beta = 1$GeV	$\beta = 1.2$GeV
$m_q = 0.2$GeV	4.6	2.6	3.6	4.2
$m_q = 0.3$GeV	3.2	5.1	2.3	3.1

IV. DISCUSSIONS AND CONCLUSIONS

In this work, we apply the PQCD method to study the branching ratio of $\overline{B}^0 \to \Lambda_c^+ \overline{p}$. The new measurements on the branching ratio by CLEO and Belle indicate that the previous theoretically estimated branching ratio in various phenomenological models are obviously larger than the data, therefore a better treatment is needed. We take the PQCD as an approach to try since it has been successful in dealing with mesonic B decays. As we discussed in the introduction, the PQCD approach should be well applicable in this case, because the momentum match demands the gluons exchanged between quarks to be "hard" and the perturbative contributions dominate. Indeed, our numerical result is satisfactorily consistent with the measured value by the Belle collaboration.

There are model-related parameters in various wavefunctions which may influence the numerical results. We obtain them by fitting data, thus the theoretical uncertainties would be much reduced. In fact, except the part involving the wavefunctions of the hadrons, all the calculations are done in the framework of PQCD. As long as the perturbative approach is applicable, the derivations follow the general principles of quantum field theory. Therefore the model-dependence is further reduced to minimum.

There is another source of possible uncertainty coming from the prescription of demanding the Sudakov factor $\exp(-s)$ to be less than one as proposed in Ref.[24]. This truncation seems to be just a convenient choice to suppress the amplitude. We therefore have carried out a calculation without the truncation for a comparison. Indeed, we find in certain kinematic regions, the Sudakov factor is larger than one. Therefore without Sudakov truncation the resulting branching ratio is larger, sometimes by a factor of two. With more accurate data, one can pin down the detailed differences. Since our results show that the predicted branching ratio with Sudakov truncation is close to experimental range, the prescription outlined in Ref.[24] is a reasonable way for calculations.

In conclusion, we have calculated the branching ratio of $\overline{B}^0 \to \Lambda_c^+ \overline{p}$ in the PQCD approach. We find that the predicted branching ratio for BR($\overline{B}^0 \to \Lambda_c^+ \overline{p}$) in the PQCD approach can vary over a range of $(2.3 \sim 5.1) \times 10^{-5}$ with the largest uncertainty coming from the parameters in the wave function of Λ_c. With the favored values for the parameters in the Λ_c^+ wave function, $\beta = 1$ GeV and $m_q = 0.3$ GeV, the branching ratio is about 2.3×10^{-5} which is satisfactorily consistent with the value measured by experiments.

Acknowledgments:

We thank Prof. Cheng who suggested us to investigate this interesting topic for fruitful discussions. This work is supported in part by the NNSF, NSC and NCTS.

Appendix A: the b measures

The ordinary b measure is defined as

$$[d\mathbf{b}] = \frac{d^2\mathbf{b}}{(2\pi)^2}. \tag{24}$$

The explicit forms of $[D\mathbf{b}]^i$ for $i-$th diagram of the first type in Fig. 1 are given by

$$\begin{aligned}
&[D\mathbf{b}]^{(a1)} = [D\mathbf{b}]^{(a2)} = [d\mathbf{b}_2][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a3)} = [D\mathbf{b}]^{(a15)} = [D\mathbf{b}]^{(a23)} = [d\mathbf{b}_2][d\mathbf{b}_3][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a4)} = [D\mathbf{b}]^{(a8)} = [D\mathbf{b}]^{(a12)} = [D\mathbf{b}]^{(a24)} = [d\mathbf{b}_1][d\mathbf{b}_2][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a5)} = [D\mathbf{b}]^{(a22)} = [D\mathbf{b}]^{(a36)} = [d\mathbf{b}_2][d\mathbf{b}_2'][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a6)} = [D\mathbf{b}]^{(a7)} = [d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a9)} = [d\mathbf{b}_1][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a10)} = [D\mathbf{b}]^{(a17)} = [d\mathbf{b}_2][d\mathbf{b}_3][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a11)} = [d\mathbf{b}_1][d\mathbf{b}_2][d\mathbf{b}_2'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a13)} = [D\mathbf{b}]^{(a14)} = [D\mathbf{b}]^{(a29)} = [D\mathbf{b}]^{(a33)} = [d\mathbf{b}_2][d\mathbf{b}_3][d\mathbf{b}_1'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a16)} = [D\mathbf{b}]^{(a19)} = [d\mathbf{b}_2][d\mathbf{b}_3][d\mathbf{b}_2'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a18)} = [d\mathbf{b}_1][d\mathbf{b}_2][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a20)} = [D\mathbf{b}]^{(a30)} = [D\mathbf{b}]^{(a34)} = [d\mathbf{b}_1][d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a21)} = [d\mathbf{b}_1][d\mathbf{b}_1'][d\mathbf{b}_2'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a25)} = [D\mathbf{b}]^{(a26)} = [d\mathbf{b}_2'][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a27)} = [D\mathbf{b}]^{(a28)} = [d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_2'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a31)} = [D\mathbf{b}]^{(a32)} = [d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_3'][d\mathbf{b}_q],\\
&[D\mathbf{b}]^{(a35)} = [d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_3'][d\mathbf{b}_q].
\end{aligned} \tag{25}$$

The explicit forms of $[D\mathbf{b}]^i$ for $i-$th diagram of the second type in Fig. 2 are given by

$$\begin{aligned}
&[D\mathbf{b}]^{(b1)} = [D\mathbf{b}]^{(a6)},\ \ [D\mathbf{b}]^{(b2)} = [D\mathbf{b}]^{(a7)},\ \ [D\mathbf{b}]^{(b3)} = [D\mathbf{b}]^{(a13)},\ \ [D\mathbf{b}]^{(b4)} = [D\mathbf{b}]^{(a14)},\\
&[D\mathbf{b}]^{(b5)} = [D\mathbf{b}]^{(a20)},\ [D\mathbf{b}]^{(b6)} = [D\mathbf{b}]^{(a21)}, [D\mathbf{b}]^{(b7)} = [D\mathbf{b}]^{(a27)},\ \ [D\mathbf{b}]^{(b8)} = [D\mathbf{b}]^{(a28)}.
\end{aligned} \tag{26}$$

The explicit forms of $[D\mathbf{b}]^i$ for $i-$th diagram of the third type in Fig. 3 are given by

$$[D\mathbf{b}]^{(c1)} = [d\mathbf{b}_2][d\mathbf{b}_1'][d\mathbf{b}_2'][d\mathbf{b}_q],\ \ [D\mathbf{b}]^{(c2)} = [d\mathbf{b}_2][d\mathbf{b}_3][d\mathbf{b}_1'][d\mathbf{b}_q]. \tag{27}$$

Appendix B: Hard scattering amplitudes $H^{i,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c})$

Expressions of amplitude $H^{i,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c})$ for each diagram in Fig. 1. There is only one Lorentz structure for the γ-matrix in the effective Hamiltonian, $O_\mu = \gamma_\mu L$.

For the hard amplitude of Fig.1(a1):

$$\begin{aligned}
&H^{a1,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) =\\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{a'c} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)\right] g_s^4\\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda(\not{q}_1 - \not{k}_2 - \not{k}_2')\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2[(q_1-k_2-k_2')^2-m_b^2][(p-k_2+k_1')^2-m_b^2]}\\
&= \frac{C_N^{a1} g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda(\not{q}_1 - \not{k}_2 - \not{k}_2')\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a1} + (\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a1} + (\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a1} + (-\mathbf{q}_T+\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][D_{a1} + (\mathbf{k}_{2T}+\mathbf{k}_{2T}'+\mathbf{k}_{3T}')^2]}
\end{aligned} \tag{28}$$

with

$$\begin{aligned}
&A_{a1} = (r^2-1)x_2x_2'm_B^2, & &B_{a1} = (r^2-1)x_3'(y-1)m_B^2,\\
&C_{a1} = m_b^2 + ((r^2-1)x_2'+1)(x_2-y)m_B^2, & &D_{a1} = m_b^2 - (r^2(x_1'-1)-x_1')(x_2-1)m_B^2
\end{aligned} \tag{29}$$

and the color factor

$$C_N^{a1} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{a'c} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{2}{3}C_1 + \frac{2}{3}C_2. \tag{30}$$

For the hard amplitude of Fig.1(a2):

$$\begin{aligned}
&H^{a2,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})=\\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{a'c}+C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right)\right]g_s^4\\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{p}-\not{k}_2+\not{k}_1'+m_b)\gamma^\theta(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2[(q-k_3')^2-m_b^2][(p-k_2+k_1')^2-m_b^2]}\\
&=\frac{C_N^{a2}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{p}-\not{k}_2+\not{k}_1'+m_b)\gamma^\theta(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a2}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a2}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a2}+\mathbf{k}_{3T}'^2][D_{a2}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}'+\mathbf{k}_{3T}')^2]}
\end{aligned} \tag{31}$$

with

$$\begin{aligned}
&A_{a2}=(r^2-1)x_2x_2'm_B^2, && B_{a2}=(r^2-1)x_3'(y-1)m_B^2,\\
&C_{a2}=m_b^2+(-x_3'r^2+x_3'-1)m_B^2, && D_{a2}=m_b^2-(r^2(x_1'-1)-x_1')(x_2-1)m_B^2
\end{aligned} \tag{32}$$

and the color factor

$$C_N^{a2}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{a'c}+C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right)=\frac{16}{3}C_1-\frac{16}{3}C_2. \tag{33}$$

For the hard amplitude of Fig.1(a3):

$$\begin{aligned}
&H^{a3,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})=\\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{a'c}+C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right)\right]g_s^4\\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2[(q-k_3')^2-m_b^2][(p-k_3+k_2')^2-m_c^2]}\\
&=\frac{C_N^{a3}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu)_{\alpha\alpha'}}{[A_{a3}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a3}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a3}+\mathbf{k}_{3T}'^2][D_{a3}+(\mathbf{k}_{3T}-\mathbf{k}_{2T}')^2]}
\end{aligned} \tag{34}$$

with

$$\begin{aligned}
&A_{a3}=(r^2-1)x_2x_2'm_B^2, && B_{a3}=(r^2-1)x_3'(y-1)m_B^2,\\
&C_{a3}=m_b^2+(-x_3'r^2+x_3'-1)m_B^2, && D_{a3}=m_c^2-(r^2(x_2'-1)-x_2')(x_3-1)m_B^2
\end{aligned} \tag{35}$$

and the color factor

$$C_N^{a3}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{a'c}+C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right)=\frac{16}{3}C_1-\frac{16}{3}C_2. \tag{36}$$

For the hard amplitude of Fig.1(a4):

$$\begin{aligned}
&H^{a4,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})=\\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^i)_{ac'}(T^j)_{ca'}+C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right)\right]g_s^4\\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')O_\mu(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(p-k_1+k_2')^2[(q-k_3')^2-m_b^2]}\\
&=\frac{C_N^{a4}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')O_\mu(\not{q}-\not{k}_3'+m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a4}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a4}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a4}+(\mathbf{k}_{1T}-\mathbf{k}_{2T}')^2][D_{a4}+\mathbf{k}_{3T}'^2]}
\end{aligned} \tag{37}$$

with

$$\begin{aligned} &A_{a4} = (r^2-1)x_2 x_2' m_B^2, && B_{a4} = (r^2-1)x_3'(y-1)m_B^2, \\ &C_{a4} = -(r^2-1)(x_1-1)x_2' m_B^2, && D_{a4} = m_b^2 + (-x_3' r^2 + x_3' - 1)m_B^2 \end{aligned} \tag{38}$$

and the color factor

$$C_N^{a4} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^i)_{ac'}(T^j)_{ca'} + C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right) = \frac{16}{3}C_1 - \frac{16}{3}C_2. \tag{39}$$

For the hard amplitude of Fig.1(a5):

$$\begin{aligned} &H^{a5,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^i)_{ac'}(T^j)_{ca'} + C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right)\right] g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q} - \not{k}_3' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu(-\not{p}' - \not{k}_2 + \not{k}_3')\gamma^\theta)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(-p'-k_2+k_3')^2[(q-k_3')^2-m_b^2]} \\ &= \frac{C_N^{a5} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q} - \not{k}_3' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu(-\not{p}' - \not{k}_2 + \not{k}_3')\gamma^\theta)_{\alpha\alpha'}}{[A_{a5} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{a5} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a5} + (\mathbf{k}_{2T} - \mathbf{k}_{3T}')^2][D_{a5} + \mathbf{k}_{3T}'^2]} \end{aligned} \tag{40}$$

with

$$\begin{aligned} &A_{a5} = (r^2-1)x_2 x_2' m_B^2, && B_{a5} = (r^2-1)x_3'(y-1)m_B^2, \\ &C_{a5} = -(r^2-1)(x_3'-1)x_2 m_B^2, && D_{a5} = m_b^2 + (-x_3' r^2 + x_3' - 1)m_B^2 \end{aligned} \tag{41}$$

and the color factor

$$C_N^{a5} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^i)_{ac'}(T^j)_{ca'} + C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right) = \frac{16}{3}C_1 - \frac{16}{3}C_2. \tag{42}$$

For the hard amplitude of Fig.1(a6):

$$\begin{aligned} &H^{a6,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ &\frac{(\gamma_\theta(\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(-p'+q_2-k_2+k_1')^2(k_2+k_2'-q_2)^2[(p-k_2+k_1')^2-m_b^2]} \\ &= \frac{C_N^{a6} g_s^4 (\gamma_\theta(\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a6} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{a6} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{a6} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}' + \mathbf{q}_T)^2][D_{a6} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2]} \end{aligned} \tag{43}$$

with

$$\begin{aligned} &A_{a6} = (r^2-1)x_2 x_2' m_B^2, && B_{a6} = (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \\ &C_{a6} = (r^2-1)(x_2+y-1)x_2' m_B^2, && D_{a6} = m_b^2 - (r^2(x_1'-1) - x_1')(x_2-1)m_B^2 \end{aligned} \tag{44}$$

and the color factor

$$C_N^{a6} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{2}{3}C_1 + \frac{2}{3}C_2. \tag{45}$$

For the hard amplitude of Fig.1(a7):

$$\begin{aligned}
&H^{a7,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x'_i, y, \mathbf{k}_T, \mathbf{k}'_T, \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^iT^j)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^iT^iT^j)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda(-\not{p}' - \not{k}_2 + \not{k}'_1)\gamma_\theta)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}'_1 + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2 + k'_2)^2(-p' + q_2 - k_2 + k'_1)^2(-p' - k_2 + k'_1)^2[(p - k_2 + k'_1)^2 - m_b^2]} \\
&= \frac{C_N^{a7} g_s^4(\gamma_\lambda(-\not{p}' - \not{k}_2 + \not{k}'_1)\gamma_\theta)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}'_1 + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a7} + (\mathbf{k}_{2T} + \mathbf{k}'_{2T})^2][B_{a7} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}'_{1T})^2][C_{a7} + (\mathbf{k}_{2T} - \mathbf{k}'_{1T})^2][D_{a7} + (\mathbf{k}_{2T} - \mathbf{k}'_{1T})^2]}
\end{aligned} \tag{46}$$

with

$$\begin{aligned}
A_{a7} &= (r^2 - 1)x_2x'_2m_B^2, & B_{a7} &= (1 - r^2)(x'_1 - 1)(x_2 + y - 1)m_B^2, \\
C_{a7} &= (1 - r^2)(x'_1 - 1)x_2m_B^2, & D_{a7} &= m_b^2 - (r^2(x'_1 - 1) - x'_1)(x_2 - 1)m_B^2
\end{aligned} \tag{47}$$

and the color factor

$$C_N^{a7} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^iT^j)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^iT^iT^j)_{cc'}\delta_{aa'}\right) = \frac{16}{3}C_1 - \frac{16}{3}C_2. \tag{48}$$

For the hard amplitude of Fig.1(a8):

$$\begin{aligned}
&H^{a8,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x'_i, y, \mathbf{k}_T, \mathbf{k}'_T, \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{q}_1 - \not{k}_2 - \not{k}'_2 + m_b)\gamma^\theta]_{\gamma\rho}(\gamma^\lambda(-\not{q}_2 + \not{k}_1 + \not{k}'_3 + m_c)O^\mu)_{\alpha\alpha'}}{(k_2 + k'_2)^2(q_2 - k_3)^2[(q_1 - k_2 - k'_2)^2 - m_b^2][(-q_2 + k_1 + k'_3)^2 - m_c^2]} \\
&= \frac{C_N^{a8} g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[O_\mu(\not{q}_1 - \not{k}_2 - \not{k}'_2 + m_b)\gamma^\theta]_{\gamma\rho}(\gamma^\lambda(-\not{q}_2 + \not{k}_1 + \not{k}'_3 + m_c)O^\mu)_{\alpha\alpha'}}{[A_{a8} + (\mathbf{k}_{2T} + \mathbf{k}'_{2T})^2][B_{a8} + (\mathbf{q}_T + \mathbf{k}'_{3T})^2][C_{a8} + (\mathbf{k}_{2T} + \mathbf{k}'_{2T} - \mathbf{q}_T)^2][D_{a8} + (\mathbf{k}_{1T} + \mathbf{k}'_{3T} + \mathbf{q}_T)^2]}
\end{aligned} \tag{49}$$

with

$$\begin{aligned}
A_{a8} &= (r^2 - 1)x_2x'_2m_B^2, & B_{a8} &= (r^2 - 1)x'_3(y - 1)m_B^2, \\
C_{a8} &= m_b^2 + ((r^2 - 1)x'_2 + 1)(x_2 - y)m_B^2, & D_{a8} &= m_c^2 + (r^2(x'_3 - 1) - x'_3)(x_1 + y - 1)m_B^2
\end{aligned} \tag{50}$$

and the color factor

$$C_N^{a8} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right) = -\frac{2}{3}C_1 + \frac{8}{3}C_2. \tag{51}$$

For the hard amplitude of Fig.1(a9):

$$\begin{aligned}
&H^{a9,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x'_i, y, \mathbf{k}_T, \mathbf{k}'_T, \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^i)_{cc'}(T^iT^j)_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(-\not{q}_2 + \not{k}_1 + \not{k}'_3 + m_c)\gamma^\theta(\not{q}_1 - \not{k}_3 - \not{k}'_1 + m_c)O^\mu)_{\alpha\alpha'}}{(k_2 + k'_2)^2(q_2 - k_3)^2[(q_1 - k_3 - k'_1)^2 - m_c^2][(-q_2 + k_1 + k'_3)^2 - m_c^2]} \\
&= \frac{C_N^{a9} g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(-\not{q}_2 + \not{k}_1 + \not{k}'_3 + m_c)\gamma^\theta(\not{q}_1 - \not{k}_3 - \not{k}'_1 + m_c)O^\mu)_{\alpha\alpha'}}{[A_{a9} + (\mathbf{k}_{2T} + \mathbf{k}'_{2T})^2][B_{a9} + (\mathbf{q}_T + \mathbf{k}'_{3T})^2][C_{a9} + (\mathbf{k}_{3T} + \mathbf{k}'_{1T} - \mathbf{q}_T)^2][D_{a9} + (\mathbf{k}_{1T} + \mathbf{k}'_{3T} + \mathbf{q}_T)^2]}
\end{aligned} \tag{52}$$

with

$$\begin{aligned} &A_{a9} = (r^2-1)x_2x'_2m_B^2, && B_{a9} = (r^2-1)x'_3(y-1)m_B^2, \\ &C_{a9} = m_c^2 + ((r^2-1)x'_1+1)(x_3-y)m_B^2, && D_{a9} = m_c^2 + (r^2(x'_3-1)-x'_3)(x_1+y-1)m_B^2 \end{aligned} \tag{53}$$

and the color factor

$$C_N^{a9} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^i)_{cc'}(T^iT^j)_{aa'}\right) = -\frac{2}{3}C_1 + \frac{8}{3}C_2. \tag{54}$$

For the hard amplitude of Fig.1(a10):

$$\begin{aligned} &H^{a10,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x'_i,y,\mathbf{k}_T,\mathbf{k}'_T,\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^i)_{cc'}(T^jT^i)_{aa'}\right)\right]g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\theta(\not p - \not k_3 + \not k'_2 + m_c)\gamma^\lambda(\not q_1 - \not k_3 - \not k'_1 + m_c)O^\mu)_{\alpha\alpha'}}{(k_2+k'_2)^2(q_2-k_3)^2[(q_1-k_3-k'_1)^2-m_c^2][(p-k_3+k'_2)^2-m_c^2]} \\ &= \frac{C_N^{a10}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\theta(\not p - \not k_3 + \not k'_2 + m_c)\gamma^\lambda(\not q_1 - \not k_3 - \not k'_1 + m_c)O^\mu)_{\alpha\alpha'}}{[A_{a10}+(\mathbf{k}_{2T}+\mathbf{k}'_{2T})^2][B_{a10}+(\mathbf{q}_T+\mathbf{k}'_{3T})^2][C_{a10}+(\mathbf{k}_{3T}+\mathbf{k}'_{1T}-\mathbf{q}_T)^2][D_{a10}+(\mathbf{k}_{3T}-\mathbf{k}'_{2T})^2]} \end{aligned} \tag{55}$$

with

$$\begin{aligned} &A_{a10} = (r^2-1)x_2x'_2m_B^2, && B_{a10} = (r^2-1)x'_3(y-1)m_B^2, \\ &C_{a10} = m_c^2 + ((r^2-1)x'_1+1)(x_3-y)m_B^2, && D_{a10} = m_c^2 - (r^2(x'_2-1)-x'_2)(x_3-1)m_B^2 \end{aligned} \tag{56}$$

and the color factor

$$C_N^{a10} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^iT^i)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^i)_{cc'}(T^jT^i)_{aa'}\right) = \frac{16}{3}C_1 + \frac{8}{3}C_2. \tag{57}$$

For the hard amplitude of Fig.1(a11):

$$\begin{aligned} &H^{a11,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x'_i,y,\mathbf{k}_T,\mathbf{k}'_T,\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^j)_{ca'}(T^iT^i)_{ac'} + C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)\right]g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not p - \not k_1 + \not k'_2)O_\mu]_{\gamma\rho}[\gamma^\lambda(-\not q_2 + \not k_1 + \not k'_3 + m_c)O^\mu)_{\alpha\alpha'}}{(k_2+k'_2)^2(q_2-k_3)^2(p-k_1+k'_2)^2[(-q2+k_1+k'_3)^2-m_c^2]} \\ &= \frac{C_N^{a11}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not p - \not k_1 + \not k'_2)O_\mu]_{\gamma\rho}[\gamma^\lambda(-\not q_2 + \not k_1 + \not k'_3 + m_c)O^\mu)_{\alpha\alpha'}}{[A_{a11}+(\mathbf{k}_{2T}+\mathbf{k}'_{2T})^2][B_{a11}+(\mathbf{q}_T+\mathbf{k}'_{3T})^2][C_{a11}+(\mathbf{k}_{1T}-\mathbf{k}'_{2T})^2][D_{a11}+(\mathbf{k}_{1T}+\mathbf{k}'_{3T}+\mathbf{q}_T)^2]} \end{aligned} \tag{58}$$

with

$$\begin{aligned} &A_{a11} = (r^2-1)x_2x'_2m_B^2, && B_{a11} = (r^2-1)x'_3(y-1)m_B^2, \\ &C_{a11} = (1-r^2)(x_1-1)x'_2m_B^2, && D_{a11} = m_c^2 + (r^2(x'_3-1)-x'_3)(x_1+y-1)m_B^2 \end{aligned} \tag{59}$$

and the color factor

$$C_N^{a11} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^j)_{ca'}(T^iT^i)_{ac'} + C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right) = \frac{16}{3}C_1 + \frac{8}{3}C_2. \tag{60}$$

For the hard amplitude of Fig.1(a12):

$$
\begin{aligned}
&H^{a12,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^j)_{ca'}(T^iT^i)_{ac'}+C_2(T^j)_{bb'}(T^i)_{cc'}(T^iT^j)_{aa'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(-\not{q}_2+\not{k}_1+\not{k}_3'+m_c)O^\mu(-\not{p}-\not{k}_2+\not{k}_3')\gamma^\theta]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3)^2(-p'-k_2+k_3')^2[(-q2+k_1+k_3')^2-m_c^2]} \\
&=\frac{C_N^{a12}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(-\not{q}_2+\not{k}_1+\not{k}_3'+m_c)O^\mu(-\not{p}-\not{k}_2+\not{k}_3')\gamma^\theta]_{\alpha\alpha'}}{[A_{a12}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a12}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a12}+(\mathbf{k}_{2T}-\mathbf{k}_{3T}')^2][D_{a12}+(\mathbf{k}_{1T}+\mathbf{k}_{3T}'+\mathbf{q}_T)^2]}
\end{aligned}
\tag{61}
$$

with

$$
\begin{aligned}
A_{a12}&=(r^2-1)x_2x_2'm_B^2, & B_{a12}&=(r^2-1)x_3'(y-1)m_B^2, \\
C_{a12}&=(1-r^2)(x_3'-1)x_2m_B^2, & D_{a12}&=m_c^2+(r^2(x_3'-1)-x_3')(x_1+y-1)m_B^2
\end{aligned}
\tag{62}
$$

and the color factor

$$
C_N^{a12}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^j)_{ca'}(T^iT^i)_{ac'}+C_2(T^j)_{bb'}(T^i)_{cc'}(T^iT^j)_{aa'}\right)=\frac{16}{3}C_1+\frac{8}{3}C_2.
\tag{63}
$$

For the hard amplitude of Fig.1(a13):

$$
\begin{aligned}
&H^{a13,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'}+C_2(T^j)_{bb'}(T^i)_{aa'}(T^jT^i)_{cc'}\right)\right]g_s^4 \\
&\frac{[\gamma_\theta(\not{k}_2+\not{k}_2'-\not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{(k_2+k_2')^2(p-q_1-k_2+k_1')^2(k_2+k_2'-q_2)^2[(q1-k_3-k_1')^2-m_c^2]} \\
&=\frac{C_N^{a13}g_s^4[\gamma_\theta(\not{k}_2+\not{k}_2'-\not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{[A_{a13}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a13}+(\mathbf{q}_T+\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][C_{a13}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}'+\mathbf{q}_T)^2][D_{a13}+(\mathbf{k}_{1T}'+\mathbf{k}_{3T}-\mathbf{q}_T)^2]}
\end{aligned}
\tag{64}
$$

with

$$
\begin{aligned}
A_{a13}&=(r^2-1)x_2x_2'm_B^2, & B_{a13}&=(1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \\
C_{a13}&=(r^2-1)(x_2+y-1)x_2'm_B^2, & D_{a13}&=m_c^2+(x_1'(r^2-1)+1)(x_3-y)m_B^2
\end{aligned}
\tag{65}
$$

and the color factor

$$
C_N^{a13}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^jT^i)_{ac'}\delta_{ca'}+C_2(T^j)_{bb'}(T^i)_{aa'}(T^jT^i)_{cc'}\right)=-\frac{2}{3}C_1+\frac{8}{3}C_2.
\tag{66}
$$

For the hard amplitude of Fig.1(a14):

$$
\begin{aligned}
&H^{a14,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^iT^j)_{ac'}\delta_{ca'}+C_2(T^j)_{bb'}(T^i)_{aa'}(T^iT^j)_{cc'}\right)\right]g_s^4 \\
&\frac{[\gamma_\lambda(-\not{p}'-\not{k}_2+\not{k}_1')\gamma_\theta]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-p'-k_2+k_1')^2(-p'-k_2+k_1')^2[(q1-k_3-k_1')^2-m_c^2]} \\
&=\frac{C_N^{a14}g_s^4[\gamma_\lambda(-\not{p}'-\not{k}_2+\not{k}_1')\gamma_\theta]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{[A_{a14}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a14}+(\mathbf{q}_T+\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][C_{a14}+(\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][D_{a14}+(\mathbf{k}_{1T}'+\mathbf{k}_{3T}-\mathbf{q}_T)^2]}
\end{aligned}
\tag{67}
$$

with

$$\begin{aligned} A_{a14} &= (r^2-1)x_2x_2'm_B^2, & B_{a14} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \\ C_{a14} &= (1-r^2)(x_1'-1)x_2m_B^2, & D_{a14} &= m_c^2+(x_1'(r^2-1)+1)(x_3-y)m_B^2 \end{aligned} \tag{68}$$

and the color factor

$$C_N^{a14} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^iT^j)_{ac'}\delta_{ca'} + C_2(T^j)_{bb'}(T^i)_{aa'}(T^iT^j)_{cc'}\right) = \frac{16}{3}C_1 + \frac{8}{3}C_2. \tag{69}$$

For the hard amplitude of Fig.1(a15):

$$\begin{aligned} &H^{a15,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)\right]g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu(\not{q}_1-\not{k}_2-\not{k}_2'+m_b)\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(k_3+k_3'-q_2)^2[(q1-k_2-k_2')^2-m_b^2]} \\ &= \frac{C_N^{a15}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu(\not{q}_1-\not{k}_2-\not{k}_2'+m_b)\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a15}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a15}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a15}+(\mathbf{k}_{3T}+\mathbf{k}_{3T}'+\mathbf{q}_T)^2][D_{a15}+(\mathbf{k}_{2T}'+\mathbf{k}_{2T}-\mathbf{q}_T)^2]} \end{aligned} \tag{70}$$

with

$$\begin{aligned} A_{a15} &= (r^2-1)x_2x_2'm_B^2, & B_{a15} &= (r^2-1)x_3'(y-1)m_B^2, \\ C_{a15} &= (r^2-1)(x_3+y-1)x_3'm_B^2, & D_{a15} &= m_b^2+(x_2'(r^2-1)+1)(x_2-y)m_B^2 \end{aligned} \tag{71}$$

and the color factor

$$C_N^{a15} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 + \frac{2}{3}C_2. \tag{72}$$

For the hard amplitude of Fig.1(a16):

$$\begin{aligned} &H^{a16,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right)\right]g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu]_{\gamma\rho}[\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(k_3+k_3'-q_2)^2[(p-k_3+k_2')^2-m_c^2]} \\ &= \frac{C_N^{a16}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu]_{\gamma\rho}[\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu]_{\alpha\alpha'}}{[A_{a16}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a16}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a16}+(\mathbf{k}_{3T}+\mathbf{k}_{3T}'+\mathbf{q}_T)^2][D_{a16}+(\mathbf{k}_{3T}-\mathbf{k}_{2T}')^2]} \end{aligned} \tag{73}$$

with

$$\begin{aligned} A_{a16} &= (r^2-1)x_2x_2'm_B^2, & B_{a16} &= (r^2-1)x_3'(y-1)m_B^2, \\ C_{a16} &= (r^2-1)(x_3+y-1)x_3'm_B^2, & D_{a16} &= m_c^2-(r^2(x_2'-1)-x_2')(x_3-1)m_B^2 \end{aligned} \tag{74}$$

and the color factor

$$C_N^{a16} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 - \frac{16}{3}C_2. \tag{75}$$

For the hard amplitude of Fig.1(a17):

$$\begin{aligned}
&H^{a17,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'}+C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)\gamma^\theta(\not{q}_1-\not{k}_1-\not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(k_3+k_3'-q_2)^2(q_1-k_1-k_1')^2} \\
&=\frac{C_N^{a17}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)\gamma^\theta(\not{q}_1-\not{k}_1-\not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a17}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a17}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a17}+(\mathbf{k}_{3T}+\mathbf{k}_{3T}'+\mathbf{q}_T)^2][D_{a17}+(\mathbf{k}_{1T}+\mathbf{k}_{1T}'-\mathbf{q}_T)^2]}
\end{aligned} \tag{76}$$

with

$$\begin{aligned}
&A_{a17}=(r^2-1)x_2x_2'm_B^2, && B_{a17}=(r^2-1)x_3'(y-1)m_B^2, \\
&C_{a17}=(r^2-1)(x_3+y-1)x_3'm_B^2, && D_{a17}=(r^2-1)(x_1'-1)(x_1-y)m_B^2
\end{aligned} \tag{77}$$

and the color factor

$$C_N^{a17}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'}+C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)=-\frac{8}{3}C_1+\frac{2}{3}C_2. \tag{78}$$

For the hard amplitude of Fig.1(a18):

$$\begin{aligned}
&H^{a18,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')\gamma^\lambda(\not{q}_1-\not{k}_1-\not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(p-k_1+k_2')^2(q_1-k_1-k_1')^2} \\
&=\frac{C_N^{a18}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')\gamma^\lambda(\not{q}_1-\not{k}_1-\not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a18}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a18}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a18}+(\mathbf{k}_{1T}-\mathbf{k}_{2T}')^2][D_{a18}+(\mathbf{k}_{1T}+\mathbf{k}_{1T}'-\mathbf{q}_T)^2]}
\end{aligned} \tag{79}$$

with

$$\begin{aligned}
&A_{a18}=(r^2-1)x_2x_2'm_B^2, && B_{a18}=(r^2-1)x_3'(y-1)m_B^2, \\
&C_{a18}=(1-r^2)(x_1-1)x_2'm_B^2, && D_{a18}=(r^2-1)(x_1'-1)(x_1-y)m_B^2
\end{aligned} \tag{80}$$

and the color factor

$$C_N^{a18}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^iT^i)_{cc'}\delta_{aa'}\right)=-\frac{8}{3}C_1-\frac{16}{3}C_2. \tag{81}$$

For the hard amplitude of Fig.1(a19):

$$\begin{aligned}
&H^{a19,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'}+C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu]_{\gamma\rho}(O^\mu(-\not{p}'-\not{k}_2+\not{k}_3')\gamma^\theta)_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(k_3+k_3'-q_2)^2(-p'-k_2+k_3')^2} \\
&=\frac{C_N^{a19}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\lambda(\not{k}_3+\not{k}_3'-\not{q}_2)O_\mu]_{\gamma\rho}(O^\mu(-\not{p}'-\not{k}_2+\not{k}_3')\gamma^\theta)_{\alpha\alpha'}}{[A_{a19}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a19}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a19}+(\mathbf{k}_{3T}+\mathbf{k}_{3T}'+\mathbf{q}_T)^2][D_{a19}+(\mathbf{k}_{2T}-\mathbf{k}_{3T}')^2]}
\end{aligned} \tag{82}$$

with

$$\begin{aligned} A_{a19} &= (r^2-1)x_2x_2' m_B^2, & B_{a19} &= (r^2-1)x_3'(y-1)m_B^2, \\ C_{a19} &= (r^2-1)(x_3+y-1)x_3' m_B^2, & D_{a19} &= (1-r^2)(x_3'-1)x_2 m_B^2 \end{aligned} \tag{83}$$

and the color factor

$$C_N^{a19} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'} + C_2(T^j)_{bb'}(T^j)_{aa'}(T^iT^i)_{cc'}\right) = -\frac{8}{3}C_1 - \frac{16}{3}C_2. \tag{84}$$

For the hard amplitude of Fig.1(a20):

$$\begin{aligned} & H^{a20,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ & \frac{[\gamma_\theta(\not k_2 + \not k_2' - \not q_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_1-p+k_2-k_1')^2(k_2+k_2'-q_2)^2(q_1-k_1-k_1')^2} \\ & = \frac{C_N^{a20} g_s^4[\gamma_\theta(\not k_2 + \not k_2' - \not q_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{[A_{a20} + (\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a20} + (\mathbf{k}_{2T}-\mathbf{k}_{1T}'+\mathbf{q}_T)^2][C_{a20} + (\mathbf{k}_{2T}+\mathbf{k}_{2T}'+\mathbf{q}_T)^2][D_{a20} + (-\mathbf{q}_T+\mathbf{k}_{1T}+\mathbf{k}_{1T}')^2]} \end{aligned} \tag{85}$$

with

$$\begin{aligned} A_{a20} &= x_2x_2'(r^2-1)m_B^2, & B_{a20} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\ C_{a20} &= x_2'(r^2-1)(x_2+y-1)m_B^2, & D_{a20} &= (r^2-1)(x_1'-1)(x_1-y)m_B^2 \end{aligned} \tag{86}$$

and the color factor

$$C_N^{a20} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 + \frac{2}{3}C_2. \tag{87}$$

For the hard amplitude of Fig.1(a21):

$$\begin{aligned} & H^{a21,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^j)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^iT^j)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ & \frac{[\gamma_\lambda(-\not p' - \not k_2 + \not k_1')\gamma_\theta]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(q_1-p+k_2-k_1')^2(-p'-k_2+k_1')^2(q_1-k_1-k_1')^2} \\ & = \frac{C_N^{a21} g_s^4[\gamma_\lambda(-\not p' - \not k_2 + \not k_1')\gamma_\theta]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{[A_{a21} + (\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a21} + (\mathbf{q}_T+\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][C_{a21} + (\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][D_{a21} + (\mathbf{k}_{1T}+\mathbf{k}_{1T}'-\mathbf{q}_T)^2]} \end{aligned} \tag{88}$$

with

$$\begin{aligned} A_{a21} &= x_2x_2'(r^2-1)m_B^2, & B_{a21} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\ C_{a21} &= (r^2-1)x_2'(x_2+y-1)m_B^2, & D_{a21} &= (r^2-1)(x_1'-1)(x_1-y)m_B^2, \end{aligned} \tag{89}$$

and the color factor

$$C_N^{a21} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^j)_{ac'}(T^i)_{ca'} + C_2(T^j)_{bb'}(T^iT^iT^j)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 - \frac{16}{3}C_2. \tag{90}$$

For the hard amplitude of Fig.1(a22):

$$\begin{aligned}
&H^{a22,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q}_1-\not{k}_2-\not{k}_2'+m_b)\gamma^\theta]_{\gamma\rho}[O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(q_2-p'+k_2')^2[(q_1-k_2-k_2')^2-m_b^2]} \\
&=\frac{C_N^{a22}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[O_\mu(\not{q}_1-\not{k}_2-\not{k}_2'+m_b)\gamma^\theta]_{\gamma\rho}[O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{[A_{a22}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a22}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a22}+(\mathbf{k}_{2T}'-\mathbf{q}_T)^2][D_{a22}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}'-\mathbf{q}_T)^2]}
\end{aligned} \tag{91}$$

with

$$\begin{aligned}
&A_{a22}=x_2x_2'(r^2-1)m_B^2, && B_{a22}=(r^2-1)x_3'(y-1)m_B^2 \\
&C_{a22}=(1-r^2)(x_2'-1)(y-1)m_B^2, && D_{a22}=m_b^2+((r^2-1)x_2'+1)(x_2-y)m_B^2,
\end{aligned} \tag{92}$$

and the color factor

$$C_N^{a22}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)=-\frac{8}{3}C_1+\frac{8}{3}C_2. \tag{93}$$

For the hard amplitude of Fig.1(a23):

$$\begin{aligned}
&H^{a23,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{aa'}(T^i)_{cc'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(q_2-p'+k_2')^2[(p-k_3+k_2')^2-m_c^2]} \\
&=\frac{C_N^{a23}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\theta(\not{p}-\not{k}_3+\not{k}_2'+m_c)O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{[A_{a23}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a23}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a23}+(\mathbf{k}_{2T}'-\mathbf{q}_T)^2][D_{a23}+(\mathbf{k}_{3T}-\mathbf{k}_{2T}')^2]}
\end{aligned} \tag{94}$$

with

$$\begin{aligned}
&A_{a23}=x_2x_2'(r^2-1)m_B^2, && B_{a23}=(r^2-1)x_3'(y-1)m_B^2 \\
&C_{a23}=(1-r^2)(x_2'-1)(y-1)m_B^2, && D_{a23}=m_c^2-(r^2(x_2'-1)-x_2')(x_3-1)m_B^2,
\end{aligned} \tag{95}$$

and the color factor

$$C_N^{a23}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{aa'}(T^i)_{cc'}\right)=-\frac{8}{3}C_1+\frac{8}{3}C_2. \tag{96}$$

For the hard amplitude of Fig.1(a24):

$$\begin{aligned}
&H^{a24,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c})= \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)\right]g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')O_\mu]_{\gamma\rho}[O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(q_2-p'+k_2')^2(p-k_1+k_2')^2} \\
&=\frac{C_N^{a24}g_s^4(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}[\gamma^\theta(\not{p}-\not{k}_1+\not{k}_2')O_\mu]_{\gamma\rho}[O^\mu(\not{q}_2-\not{p}'+\not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{[A_{a24}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a24}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a24}+(\mathbf{k}_{2T}'-\mathbf{q}_T)^2][D_{a24}+(\mathbf{k}_{1T}-\mathbf{k}_{2T}')^2]}
\end{aligned} \tag{97}$$

with

$$\begin{array}{ll} A_{a24} = x_2 x_2'(r^2-1)m_B^2, & B_{a24} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a24} = (1-r^2)(x_2'-1)(y-1)m_B^2, & D_{a24} = (1-r^2)(x_1-1)x_2' m_B^2, \end{array} \tag{98}$$

and the color factor

$$C_N^{a24} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1 (T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'} + C_2 (T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{99}$$

For the hard amplitude of Fig.1(a25):

$$\begin{aligned} & H^{a25,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1 (T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'} + C_2 (T^j)_{bb'}(T^jT^i)_{aa'}(T^i)_{cc'}\right)\right] g_s^4 \\ & \frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{q}_2 - \not{p}' - \not{k}_2)\gamma^\theta(\not{q}_2 - \not{p}' + \not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(q_2-p'+k_2')^2(q_2-p'-k_2)^2} \\ & = \frac{C_N^{a25} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{q}_2 - \not{p}' - \not{k}_2)\gamma^\theta(\not{q}_2 - \not{p}' + \not{k}_2')\gamma^\lambda]_{\alpha\alpha'}}{[A_{a25} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{a25} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a25} + (\mathbf{k}_{2T}' - \mathbf{q}_T)^2][D_{a25} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2]} \end{aligned} \tag{100}$$

with

$$\begin{array}{ll} A_{a25} = x_2 x_2'(r^2-1)m_B^2, & B_{a25} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a25} = (1-r^2)(x_2'-1)(y-1)m_B^2, & D_{a25} = (r^2-1)(x_2+y-1)m_B^2, \end{array} \tag{101}$$

and the color factor

$$C_N^{a25} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1 (T^j)_{bb'}(T^i)_{ac'}(T^jT^i)_{ca'} + C_2 (T^j)_{bb'}(T^jT^i)_{aa'}(T^i)_{cc'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{102}$$

For the hard amplitude of Fig.1(a26):

$$\begin{aligned} & H^{a26,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1 (T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'} + C_2 (T^j)_{bb'}(T^iT^j)_{aa'}(T^i)_{cc'}\right)\right] g_s^4 \\ & \frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{q}_2 - \not{p}' - \not{k}_2)\gamma^\lambda(-\not{p}' - \not{k}_2 + \not{k}_3')\gamma^\theta]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-k_3')^2(q_2-p'+k_2')^2(-p'-k_2+k_3')^2} \\ & = \frac{C_N^{a26} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}(\gamma_\theta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{q}_2 - \not{p}' - \not{k}_2)\gamma^\lambda(-\not{p}' - \not{k}_2 + \not{k}_3')\gamma^\theta]_{\alpha\alpha'}}{[A_{a26} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{a26} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a26} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2][D_{a26} + (\mathbf{k}_{2T} - \mathbf{k}_{3T})^2]} \end{aligned} \tag{103}$$

with

$$\begin{array}{ll} A_{a26} = x_2 x_2'(r^2-1)m_B^2, & B_{a26} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a26} = (r^2-1)(x_2+y-1)m_B^2, & D_{a26} = (1-r^2)x_2(x_3'-1)m_B^2, \end{array} \tag{104}$$

and the color factor

$$C_N^{a26} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1 (T^j)_{bb'}(T^i)_{ac'}(T^iT^j)_{ca'} + C_2 (T^j)_{bb'}(T^iT^j)_{aa'}(T^i)_{cc'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{105}$$

For the hard amplitude of Fig.1(a27):

$$\begin{aligned}
&H^{a27,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\
&\Big[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)\Big]\,g_s^4 \\
&\frac{[\gamma_\theta(\not{k}_2+\not{k}_2'-\not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O^\mu)_{\gamma\rho}[O^\mu(-\not{p}'+\not{q}_2-\not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-p'-k_2+k_1')^2(q_2-p'-k_2)^2(-q_2+k_2+k_2')^2} \\
&=\frac{C_N^{a27}g_s^4[\gamma_\theta(\not{k}_2+\not{k}_2'-\not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O^\mu)_{\gamma\rho}[O^\mu(-\not{p}'+\not{q}_2-\not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{[A_{a27}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a27}+(\mathbf{q}_T+\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][C_{a27}+(\mathbf{k}_{2T}+\mathbf{q}_T)^2][D_{a27}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}'+\mathbf{q}_T)^2]}
\end{aligned} \tag{106}$$

with

$$\begin{aligned}
A_{a27}&=x_2x_2'(r^2-1)m_B^2, & B_{a27}&=(1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\
C_{a27}&=(r^2-1)(x_2+y-1)m_B^2, & D_{a27}&=(r^2-1)x_2'(x_2+y-1)m_B^2,
\end{aligned} \tag{107}$$

and the color factor

$$C_N^{a27}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^jT^i)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^jT^i)_{cc'}(T^i)_{aa'}\right)=-\frac{8}{3}C_1+\frac{8}{3}C_2. \tag{108}$$

For the hard amplitude of Fig.1(a28):

$$\begin{aligned}
&H^{a28,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\
&\Big[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^j)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^iT^j)_{cc'}(T^i)_{aa'}\right)\Big]\,g_s^4 \\
&\frac{[\gamma_\lambda(-\not{p}'-\not{k}_2+\not{k}_1')\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O^\mu)_{\gamma\rho}[O^\mu(-\not{p}'+\not{q}_2-\not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{(k_2+k_2')^2(q_2-p'-k_2+k_1')^2(q_2-p'-k_2)^2(-p'-k_2+k_1')^2} \\
&=\frac{C_N^{a28}g_s^4[\gamma_\lambda(-\not{p}'-\not{k}_2+\not{k}_1')\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O^\mu)_{\gamma\rho}[O^\mu(-\not{p}'+\not{q}_2-\not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{[A_{a28}+(\mathbf{k}_{2T}+\mathbf{k}_{2T}')^2][B_{a28}+(\mathbf{q}_T+\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2][C_{a28}+(\mathbf{k}_{2T}+\mathbf{q}_T)^2][D_{a28}+(\mathbf{k}_{2T}-\mathbf{k}_{1T}')^2]}
\end{aligned} \tag{109}$$

with

$$\begin{aligned}
A_{a28}&=x_2x_2'(r^2-1)m_B^2, & B_{a28}&=(1-r^2)(x_1'-1)(-1+x_2+y)m_B^2 \\
C_{a28}&=(r^2-1)(-1+x_2+y)m_B^2, & D_{a28}&=(1-r^2)(x_1'-1)x_2m_B^2,
\end{aligned} \tag{110}$$

and the color factor

$$C_N^{a28}=\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^j)_{bb'}(T^iT^j)_{ac'}(T^i)_{ca'}+C_2(T^j)_{bb'}(T^iT^j)_{cc'}(T^i)_{aa'}\right)=-\frac{8}{3}C_1+\frac{8}{3}C_2. \tag{111}$$

For the hard amplitude of Fig.1(a29):

$$\begin{aligned}
&H^{a29,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i,x_i',y,\mathbf{k}_T,\mathbf{k}_T',\mathbf{q}_T,m_B,m_{\Lambda_c}) = \\
&\Big[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^jT^i)_{ac'}\delta_{ca'}+C_2(T^jT^i)_{bb'}(T^j)_{aa'}(T^i)_{cc'}\right)\Big]\,g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p}-\not{q}_1+\not{k}_1')\gamma^\lambda]_{\beta\beta'}(O^\mu)_{\gamma\rho}[\gamma^\theta(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{(p-q_1-k_2+k_1')^2(q_2-k_3')^2(p-q_1+k_1')^2[(q_1-k_3-k_1')^2-m_c^2} \\
&=\frac{C_N^{a29}g_s^4(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p}-\not{q}_1+\not{k}_1')\gamma^\lambda]_{\beta\beta'}(O^\mu)_{\gamma\rho}[\gamma^\theta(\not{q}_1-\not{k}_3-\not{k}_1'+m_c)O^\mu]_{\alpha\alpha'}}{[A_{a29}+(\mathbf{k}_{2T}-\mathbf{k}_{1T}'+\mathbf{q}_T)^2][B_{a29}+(\mathbf{q}_T+\mathbf{k}_{3T}')^2][C_{a29}+(\mathbf{k}_{1T}'-\mathbf{q}_T)^2][D_{a29}+(\mathbf{k}_{3T}+\mathbf{k}_{1T}'-\mathbf{q}_T)^2]}
\end{aligned} \tag{112}$$

with

$$\begin{aligned} A_{a29} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \quad B_{a29} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a29} &= (1-r^2)(x_1'-1)(-1+y)m_B^2, \qquad D_{a29} = m_c^2 + (1+(r^2-1)x_1')(x_3-y)m_B^2, \end{aligned} \tag{113}$$

and the color factor

$$C_N^{a29} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^jT^i)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^j)_{aa'}(T^i)_{cc'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{114}$$

For the hard amplitude of Fig.1(a30):

$$\begin{aligned} &H^{a30,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^i)_{ac'}(T^j)_{ca'} + C_2(T^jT^i)_{bb'}(T^jT^i)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p} - \not{q}_1 + \not{k}_1')\gamma^\lambda]_{\beta\beta'}[\gamma^\theta(\not{q}_1 - \not{k}_1 - \not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(p-q_1-k_2+k_1')^2(q_2-k_3')^2(p-q_1+k_1')^2(q_1-k_1-k_1')^2} \\ &= \frac{C_N^{a30}g_s^4(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p} - \not{q}_1 + \not{k}_1')\gamma^\lambda]_{\beta\beta'}[\gamma^\theta(\not{q}_1 - \not{k}_1 - \not{k}_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a30} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][B_{a30} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a30} + (\mathbf{k}_{1T}' - \mathbf{q}_T)^2][D_{a30} + (\mathbf{k}_{1T} + \mathbf{k}_{1T}' - \mathbf{q}_T)^2]} \end{aligned} \tag{115}$$

with

$$\begin{aligned} A_{a30} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \quad B_{a30} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a30} &= (1-r^2)(x_1'-1)(-1+y)m_B^2, \qquad D_{a30} = (r^2-1)(x_1'-1)(x_1-y)m_B^2, \end{aligned} \tag{116}$$

and the color factor

$$C_N^{a30} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^i)_{ac'}(T^j)_{ca'} + C_2(T^jT^i)_{bb'}(T^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{117}$$

For the hard amplitude of Fig.1(a31):

$$\begin{aligned} &H^{a31,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ &\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^jT^i)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^jT^i)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ &\frac{(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p} - \not{q}_1 + \not{k}_1')\gamma^\lambda]_{\beta\beta'}[O_\mu\gamma^\theta(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(p-q_1-k_2+k_1')^2(q_2-k_3')^2(p-q_1+k_1')^2[(p-k_2+k_1')^2 - m_b^2]} \\ &= \frac{C_N^{a31}g_s^4(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not{p} - \not{q}_1 + \not{k}_1')\gamma^\lambda]_{\beta\beta'}[O_\mu\gamma^\theta(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a31} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][B_{a31} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a31} + (\mathbf{k}_{1T}' - \mathbf{q}_T)^2][D_{a31} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2]} \end{aligned} \tag{118}$$

with

$$\begin{aligned} A_{a31} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, \quad B_{a31} = (r^2-1)x_3'(y-1)m_B^2 \\ C_{a31} &= (1-r^2)(x_1'-1)(-1+y)m_B^2, \qquad D_{a31} = (1+x_1'(-1+x_2)+r^2(-1+x_1'+x_2-x_1'x_2))m_B^2, \end{aligned} \tag{119}$$

and the color factor

$$C_N^{a31} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^jT^i)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^jT^i)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{120}$$

For the hard amplitude of Fig.1(a32):

$$\begin{aligned}
&H^{a32,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^i)_{ac'}(T^j)_{ca'} + C_2(T^jT^i)_{bb'}(T^i)_{cc'}(T^j)_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not q_2 - \not p' + \not k_1')\gamma^\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not p - \not q_1 - \not k_2)\gamma^\theta]_{\alpha\alpha'}}{(p-q_1-k_2+k_1')^2(q_2-k_3')^2(p-q_1-k_2)^2(q_2-p'+k_1')^2} \\
&= \frac{C_N^{a32} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}[\gamma_\theta(\not q_2 - \not p' + \not k_1')\gamma^\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not p - \not q_1 - \not k_2)\gamma^\theta]_{\alpha\alpha'}}{[A_{a32} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][B_{a32} + (\mathbf{q}_T + \mathbf{k}_{3T}')^2][C_{a32} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2][D_{a32} + (-\mathbf{q}_T + \mathbf{k}_{1T}')^2]}
\end{aligned} \tag{121}$$

with

$$\begin{aligned}
A_{a32} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2, & B_{a32} &= (r^2-1)x_3'(y-1)m_B^2 \\
C_{a32} &= (r^2-1)(x_2+y-1)m_B^2, & D_{a32} &= (1-r^2)(x_1'-1)(y-1)m_B^2,
\end{aligned} \tag{122}$$

and the color factor

$$C_N^{a32} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^i)_{ac'}(T^j)_{ca'} + C_2(T^jT^i)_{bb'}(T^i)_{cc'}(T^j)_{aa'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{123}$$

For the hard amplitude of Fig.1(a33):

$$\begin{aligned}
&H^{a33,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^iT^j)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^j)_{cc'}(T^i)_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not q_2 + \not k_2 + \not k_3')\gamma_\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not q_1 - \not k_3 - \not k_1' + m_c)O^\mu]_{\alpha\alpha'}}{(q_2-k_3')^2(q_2-p'-k_2+k_1')^2(-q_2+k_2+k_3')^2[(q_1-k_3-k_1')^2-m_c^2]} \\
&= \frac{C_N^{a33} g_s^4 (\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not q_2 + \not k_2 + \not k_3')\gamma_\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma^\lambda(\not q_1 - \not k_3 - \not k_1' + m_c)O^\mu]_{\alpha\alpha'}}{[A_{a33} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][B_{a33} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][C_{a33} + (\mathbf{k}_{2T} + \mathbf{k}_{3T}' + \mathbf{q}_T)^2][D_{a33} + (-\mathbf{q}_T + \mathbf{k}_{1T}' + \mathbf{k}_{3T})^2]}
\end{aligned} \tag{124}$$

with

$$\begin{aligned}
A_{a33} &= (r^2-1)x_3'(y-1)m_B^2, & B_{a33} &= (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\
C_{a33} &= (r^2-1)(x_2+y-1)x_3'm_B^2, & D_{a33} &= m_c^2 + (1+(r^2-1)x_1')(x_3-y)m_B^2,
\end{aligned} \tag{125}$$

and the color factor

$$C_N^{a33} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^iT^j)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^j)_{cc'}(T^i)_{aa'}\right) = \frac{10}{3}C_1 + \frac{8}{3}C_2. \tag{126}$$

For the hard amplitude of Fig.1(a34):

$$\begin{aligned}
&H^{a34,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&\left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^j)_{ac'}(T^i)_{ca'} + C_2(T^jT^i)_{bb'}(T^iT^j)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not q_2 + \not k_2 + \not k_3')\gamma_\lambda]_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(q_2-k_3')^2(p-q_1-k_2+k_1')^2(-q_2+k_2+k_3')^2(q_1-k_1-k_1')^2} \\
&= \frac{C_N^{a34} g_s^4 (\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not q_2 + \not k_2 + \not k_3')\gamma_\lambda]_{\beta\beta'}[\gamma^\lambda(\not q_1 - \not k_1 - \not k_1')O_\mu]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a34} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][B_{a34} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][C_{a34} + (\mathbf{k}_{2T} + \mathbf{k}_{3T}' + \mathbf{q}_T)^2][D_{a34} + (-\mathbf{q}_T + \mathbf{k}_{1T}' + \mathbf{k}_{1T})^2]}
\end{aligned} \tag{127}$$

with

$$\begin{array}{ll} A_{a34} = (r^2-1)x_3'(y-1)m_B^2, & B_{a34} = (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\ C_{a34} = (r^2-1)(x_2+y-1)x_3'm_B^2, & D_{a34} = (r^2-1)(x_1'-1)(x_1-y)m_B^2, \end{array} \tag{128}$$

and the color factor

$$C_N^{a34} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^j)_{ac'}(T^i)_{ca'} + C_2(T^jT^i)_{bb'}(T^iT^j)_{cc'}\delta_{aa'}\right) = -\frac{8}{3}C_1 - \frac{10}{3}C_2. \tag{129}$$

For the hard amplitude of Fig.1(a35):

$$\begin{aligned} & H^{a35,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^iT^j)_{ac'}\delta_{ca'} + C_2(T^jT^i)_{bb'}(T^iT^j)_{cc'}\delta_{aa'}\right)\right] g_s^4 \\ & \frac{(\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not{q}_2 + \not{k}_2 + \not{k}_3')\gamma_\lambda]_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(q_2 - k_3')^2(q_2 - p' - k_2 + k_1')^2(-q_2 + k_2 + k_3')^2[(p - k_2 + k_1')^2 - m_b^2]} \\ & = \frac{C_N^{a35} g_s^4 (\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not{q}_2 + \not{k}_2 + \not{k}_3')\gamma_\lambda]_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{a35} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][B_{a35} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][C_{a35} + (\mathbf{k}_{2T} + \mathbf{k}_{3T}' + \mathbf{q}_T)^2][D_{a35} + (-\mathbf{k}_{1T}' + \mathbf{k}_{2T})^2]} \end{aligned} \tag{130}$$

with

$$\begin{array}{ll} A_{a35} = (r^2-1)x_3'(y-1)m_B^2, & B_{a35} = (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\ C_{a35} = (r^2-1)(x_2+y-1)x_3'm_B^2, & D_{a35} = (1 + x_1'(-1+x_2) + r^2(-1 + x_1' + x_2 - x_1'x_2))m_B^2, \end{array} \tag{131}$$

and the color factor

$$C_N^{a35} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^j)_{ac'}(T^i)_{ca'} + C_2(T^jT^i)_{bb'}(T^iT^j)_{cc'}\delta_{aa'}\right) = \frac{10}{3}C_1 - \frac{10}{3}C_2. \tag{132}$$

For the hard amplitude of Fig.1(a36):

$$\begin{aligned} & H^{a36,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\ & \left[\varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^j)_{ac'}(T^i)_{ca'} + C_2(T^jT^i)_{bb'}(T^j)_{cc'}(T^i)_{aa'}\right)\right] g_s^4 \\ & \frac{(\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not{q}_2 + \not{k}_2 + \not{k}_3')\gamma_\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{p} - \not{q}_1 - \not{k}_2 + m_b)\gamma^\lambda]_{\alpha\alpha'}}{(q_2 - k_3')^2(-p + q_1 + k_2 - k_1')^2(-q_2 + k_2 + k_3')^2(p - q_1 - k_2)^2} \\ & = \frac{C_N^{a36} g_s^4 (\gamma_\theta)_{\rho'\gamma'}[\gamma^\theta(-\not{q}_2 + \not{k}_2 + \not{k}_3')\gamma_\lambda]_{\beta\beta'}(O_\mu)_{\gamma\rho}[O^\mu(\not{p} - \not{q}_1 - \not{k}_2 + m_b)\gamma^\lambda]_{\alpha\alpha'}}{[A_{a36} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][B_{a36} + (-\mathbf{k}_{2T} + \mathbf{k}_{1T}' - \mathbf{q}_T)^2][C_{a36} + (\mathbf{k}_{2T} + \mathbf{k}_{3T}' + \mathbf{q}_T)^2][D_{a36} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2]} \end{aligned} \tag{133}$$

with

$$\begin{array}{ll} A_{a36} = (r^2-1)x_3'(y-1)m_B^2, & B_{a36} = (1-r^2)(x_1'-1)(x_2+y-1)m_B^2 \\ C_{a36} = (r^2-1)(x_2+y-1)x_3'm_B^2, & D_{a36} = (r^2-1)(x_2+y-1)m_B^2, \end{array} \tag{134}$$

and the color factor

$$C_N^{a36} = \varepsilon^{abc}\varepsilon^{a'b'c'}\left(C_1(T^jT^i)_{bb'}(T^j)_{ac'}(T^i)_{ca'} + C_2(T^jT^i)_{bb'}(T^j)_{cc'}(T^i)_{aa'}\right) = -\frac{8}{3}C_1 + \frac{8}{3}C_2. \tag{135}$$

For the hard amplitude of Fig.2(b1):

$$\begin{aligned} &H^{b1,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a6,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ &= \frac{C_N^{b1} g_s^4 (\gamma_\theta (\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda)_{\rho'\gamma'} (\gamma^\theta)_{\beta\beta'} [O_\mu (\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho} (O^\mu)_{\alpha\alpha'}}{[A_{b1} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b1} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b1} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}' + \mathbf{q}_T)^2][D_{b1} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2]} \end{aligned} \tag{136}$$

with

$$\begin{aligned} A_{b1} &= x_2 x_2' (r^2 - 1) m_B^2, & B_{b1} &= x_2(-(x_1' - 1) r^2 + x_1' - y) m_B^2, \\ C_{b1} &= x_2((r^2 - 1) x_2' - y + 1) m_B^2, & D_{b1} &= m_b^2 - (r^2 (x_1' - 1) - x_1')(x_2 - 1) m_B^2 \end{aligned} \tag{137}$$

and the color factor

$$C_N^{b1} = C_N^{a6}. \tag{138}$$

For the hard amplitude of Fig.2(b2):

$$\begin{aligned} &H^{b2,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a7,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ &= \frac{C_N^{b2} g_s^4 (\gamma_\lambda (-\not{p}' - \not{k}_2 + \not{k}_1')\gamma_\theta)_{\rho'\gamma'} (\gamma^\theta)_{\beta\beta'} [O_\mu (\not{p} - \not{k}_2 + \not{k}_1' + m_b)\gamma^\lambda]_{\gamma\rho} (O^\mu)_{\alpha\alpha'}}{[A_{b2} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b2} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b2} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][D_{b2} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2]} \end{aligned} \tag{139}$$

with

$$\begin{aligned} A_{b2} &= x_2 x_2' (r^2 - 1) m_B^2, & B_{b2} &= x_2(-(x_1' - 1) r^2 + x_1' - y) m_B^2, \\ C_{b2} &= -x_2 (r^2 - 1)(x_1' - 1) m_B^2, & D_{b2} &= m_b^2 - (x_2 - 1)(r^2 (x_1' - 1) - x_1') m_B^2 \end{aligned} \tag{140}$$

and the color factor

$$C_N^{b2} = C_N^{a7}. \tag{141}$$

For the hard amplitude of Fig.2(b3):

$$\begin{aligned} &H^{b3,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a13,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ &= \frac{C_N^{b3} g_s^4 [\gamma_\theta (\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda]_{\rho'\gamma'} (\gamma^\theta)_{\beta\beta'} (O_\mu)_{\gamma\rho} [\gamma^\lambda (\not{q}_1 - \not{k}_3 - \not{k}_1' + m_c) O^\mu]_{\alpha\alpha'}}{[A_{b3} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b3} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b3} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}' + \mathbf{q}_T)^2][D_{b3} + (\mathbf{k}_{1T}' + \mathbf{k}_{3T} - \mathbf{q}_T)^2]} \end{aligned} \tag{142}$$

with

$$\begin{aligned} A_{b3} &= x_2 x_2' (r^2 - 1) m_B^2, & B_{b3} &= x_2(-(x_1' - 1) r^2 + x_1' - y) m_B^2, \\ C_{b3} &= x_2((r^2 - 1) x_2' - y + 1) m_B^2, & D_{b3} &= m_c^2 + (x_3 - 1)((r^2 - 1) x_1' + y) m_B^2 \end{aligned} \tag{143}$$

and the color factor

$$C_N^{b3} = C_N^{a13}. \tag{144}$$

For the hard amplitude of Fig.2(b4):

$$\begin{aligned} &H^{b4,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a14,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ &= \frac{C_N^{b4} g_s^4 [\gamma_\lambda (-\not{p}' - \not{k}_2 + \not{k}_1')\gamma_\theta]_{\rho'\gamma'} (\gamma^\theta)_{\beta\beta'} (O_\mu)_{\gamma\rho} [\gamma^\lambda (\not{q}_1 - \not{k}_3 - \not{k}_1' + m_c) O^\mu]_{\alpha\alpha'}}{[A_{b4} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b4} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b4} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][D_{b4} + (\mathbf{k}_{1T}' + \mathbf{k}_{3T} - \mathbf{q}_T)^2]} \end{aligned} \tag{145}$$

with

$$\begin{array}{ll} A_{b4} = x_2 x_2'(r^2-1)m_B^2, & B_{b4} = x_2(-(x_1'-1)r^2 + x_1' - y)m_B^2, \\ C_{b4} = -x_2(r^2-1)(x_1'-1)m_B^2, & D_{b4} = m_c^2 + (x_3-1)((r^2-1)x_1' + y)m_B^2 \end{array} \tag{146}$$

and the color factor

$$C_N^{b4} = C_N^{a14}. \tag{147}$$

For the hard amplitude of Fig.2(b5):

$$\begin{aligned} & H^{b5,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a20,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ & = \frac{C_N^{b5} g_s^4 [\gamma_\theta(\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not{q}_1 - \not{k}_1 - \not{k}_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{[A_{b5} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b5} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}' + \mathbf{q}_T)^2][C_{b5} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}' + \mathbf{q}_T)^2][D_{b5} + (-\mathbf{q}_T + \mathbf{k}_{1T} + \mathbf{k}_{1T}')^2]} \end{aligned} \tag{148}$$

with

$$\begin{array}{ll} A_{b5} = x_2 x_2'(r^2-1)m_B^2, & B_{b5} = m_B^2(r^2(x_2 - x_1'x_2) + x_2(x_1' - y)), \\ C_{b5} = m_B^2 x_2(1 + (-1+r^2)x_2' - y), & D_{b5} = m_B^2(r^2(1 - x_1 + (-1+x_1)x_1') - (-1+x_1)(x_1' - y)), \end{array} \tag{149}$$

and the color factor

$$C_N^{b5} = C_N^{a20}. \tag{150}$$

For the hard amplitude of Fig.2(b6):

$$\begin{aligned} & H^{b6,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a21,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ & = \frac{C_N^{b6} g_s^4 [\gamma_\lambda(-\not{p}' - \not{k}_2 + \not{k}_1')\gamma_\theta]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}[\gamma^\lambda(\not{q}_1 - \not{k}_1 - \not{k}_1')O^\mu]_{\gamma\rho}(O_\mu)_{\alpha\alpha'}}{[A_{b6} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b6} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b6} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][D_{b6} + (\mathbf{k}_{1T} + \mathbf{k}_{1T}' - \mathbf{q}_T)^2]} \end{aligned} \tag{151}$$

with

$$\begin{array}{ll} A_{b6} = x_2 x_2'(r^2-1)m_B^2, & B_{b6} = m_B^2(r^2(x_2 - x_1'x_2) + x_2(x_1' - y)) \\ C_{b6} = -m_B^2 x_2(-1+r^2)(-1+x_1'), & D_{b6} = m_B^2(r^2(1 - x_1 + (-1+x_1)x_1') - (-1+x_1)(x_1' - y)), \end{array} \tag{152}$$

and the color factor

$$C_N^{b6} = C_N^{a21}. \tag{153}$$

For the hard amplitude of Fig.2(b7):

$$\begin{aligned} & H^{b7,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a27,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\ & = \frac{C_N^{b7} g_s^4 [\gamma_\theta(\not{k}_2 + \not{k}_2' - \not{q}_2)\gamma_\lambda]_{\rho'\gamma'}(\gamma^\theta)_{\beta\beta'}(O^\mu)_{\gamma\rho}[O^\mu(-\not{p}' + \not{q}_2 - \not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{[A_{b7} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b7} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b7} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2][D_{b7} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}' + \mathbf{q}_T)^2]} \end{aligned} \tag{154}$$

with

$$\begin{array}{ll} A_{b7} = x_2 x_2'(r^2-1)m_B^2, & B_{b7} = m_B^2 x_2(-r^2(-1+x_1') + x_1' - y) \\ C_{b7} = m_B^2 x_2(-y + r^2), & D_{b7} = m_B^2 x_2(1 + x_2'(-1+r^2) - y), \end{array} \tag{155}$$

and the color factor

$$C_N^{b7} = C_N^{a27}. \tag{156}$$

For the hard amplitude of Fig.2(b8):

$$\begin{aligned}
&H^{b8,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = H^{a28,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) \\
&= \frac{C_N^{a28} g_s^4 [\gamma_\lambda(-\not{p}' - \not{k}_2 + \not{k}_1')\gamma_\lambda]_{\rho'\gamma'} (\gamma^\theta)_{\beta\beta'} (O^\mu)_{\gamma\rho} [O^\mu(-\not{p}' + \not{q}_2 - \not{k}_2)\gamma_\lambda]_{\alpha\alpha'}}{[A_{b8} + (\mathbf{k}_{2T} + \mathbf{k}_{2T}')^2][B_{b8} + (\mathbf{q}_T + \mathbf{k}_{2T} - \mathbf{k}_{1T}')^2][C_{b8} + (\mathbf{k}_{2T} + \mathbf{q}_T)^2][D_{b8} + (\mathbf{k}_{2T} - \mathbf{k}_{1T}')^2]}
\end{aligned} \tag{157}$$

with

$$\begin{aligned}
A_{b8} &= x_2 x_2'(r^2-1)m_B^2, \quad B_{b8} = m_B^2 x_2(-r^2(-1+x_1') + x_1' - y) \\
C_{b8} &= m_B^2 x_2(-y + r^2), \quad D_{b8} = -m_B^2 x_2(-1+r^2)(-1+x_1'),
\end{aligned} \tag{158}$$

and the color factor

$$C_N^{b8} = C_N^{a28}. \tag{159}$$

For the hard amplitude of Fig.3(c1):

$$\begin{aligned}
&H^{c1,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&i\left[\varepsilon^{abc}\varepsilon^{a'b'c'} f_{jil}\left(C_1(T^jT^i)_{ac'}(T^l)_{bb'}\delta_{ca'} + C_2(T^jT^i)_{cc'}(T^l)_{bb'}\delta_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\eta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 - \not{k}_1' + m_b)\gamma_\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{(k_2+k_2')^2(-q_2+k_3')^2(p-q_1-k_2-k_1')^2[(p-k_2-k_1')^2 - m_b^2]} \\
&[g^{\theta\lambda}(p-q_1-k_2-k_1'-(-q_2+k_3'))^\eta + g^{\lambda\eta}(-q_2+k_3'-(k_2+k_2'))^\theta + g^{\eta\theta}(k_2+k_2'-(p-k_2-k_1'))^\lambda] \\
&= \frac{iC_N^{c1} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}(\gamma_\eta)_{\beta\beta'}[O_\mu(\not{p} - \not{k}_2 - \not{k}_1' + m_b)\gamma_\theta]_{\gamma\rho}(O^\mu)_{\alpha\alpha'}}{[A_{c1} + (\mathbf{k}_{2T}' + \mathbf{k}_{2T}')^2][B_{c1} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][C_{c1} + (\mathbf{k}_{2T} + \mathbf{k}_{1T}' + \mathbf{q}_T)^2][D_{c1} + (\mathbf{k}_{2T} + \mathbf{k}_{1T}')^2]} \\
&[g^{\theta\lambda}(p-q_1-k_2-k_1'-(-q_2+k_3'))^\eta + g^{\lambda\eta}(-q_2+k_3'-(k_2+k_2'))^\theta + g^{\eta\theta}(k_2+k_2'-(p-k_2-k_1'))^\lambda]
\end{aligned} \tag{160}$$

with

$$\begin{aligned}
A_{c1} &= x_2 x_2'(r^2-1)m_B^2, & B_{c1} &= (r^2-1)x_3'(y-1)m_B^2 \\
C_{c1} &= (r^2-1)(x_2+y-1)(x_1'+1)m_B^2, & D_{c1} &= m_b^2 + (r^2(x_1'+1) - x_1')(x_2-1)m_B^2,
\end{aligned} \tag{161}$$

and the color factor

$$C_N^{c1} = \varepsilon^{abc}\varepsilon^{a'b'c'} f_{jil}\left(C_1(T^jT^i)_{ac'}(T^l)_{bb'}\delta_{ca'} + C_2(T^jT^i)_{cc'}(T^l)_{bb'}\delta_{aa'}\right) = 6iC_1 - 6iC_2. \tag{162}$$

For the hard amplitude of Fig.3(c2):

$$\begin{aligned}
&H^{c2,\alpha'\beta'\gamma'\rho'\alpha\beta\gamma\rho}(x_i, x_i', y, \mathbf{k}_T, \mathbf{k}_T', \mathbf{q}_T, m_B, m_{\Lambda_c}) = \\
&i\left[\varepsilon^{abc}\varepsilon^{a'b'c'} f_{jil}\left(C_1(T^jT^i)_{ac'}(T^l)_{bb'}\delta_{ca'} + C_2(T^i)_{cc'}(T^l)_{bb'}(T^j)_{aa'}\right)\right] g_s^4 \\
&\frac{(\gamma_\lambda)_{\rho'\gamma'}(\gamma_\eta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma_\theta(\not{q}_1 - \not{k}_3 - \not{k}_1' + m_c)O^\mu]_{\alpha\alpha'}}{(k_2+k_2')^2(-q_2+k_3')^2(p-q_1-k_2-k_1')^2[(q_1-k_3-k_1')^2 - m_c^2]} \\
&[g^{\theta\lambda}(p-q_1-k_2-k_1'-(-q_2+k_3'))^\eta + g^{\lambda\eta}(-q_2+k_3'-(k_2+k_2'))^\theta + g^{\eta\theta}(k_2+k_2'-(p-k_2-k_1'))^\lambda] \\
&= \frac{iC_N^{c2} g_s^4 (\gamma_\lambda)_{\rho'\gamma'}(\gamma_\eta)_{\beta\beta'}(O_\mu)_{\gamma\rho}[\gamma_\theta(\not{q}_1 - \not{k}_3 - \not{k}_1' + m_c)O^\mu]_{\alpha\alpha'}}{[A_{c2} + (\mathbf{k}_{2T}' + \mathbf{k}_{2T}')^2][B_{c2} + (\mathbf{k}_{3T}' + \mathbf{q}_T)^2][C_{c2} + (\mathbf{k}_{2T} + \mathbf{k}_{1T}' + \mathbf{q}_T)^2][D_{c2} + (\mathbf{k}_{3T} + \mathbf{k}_{1T}' - \mathbf{q}_T)^2]} \\
&[g^{\theta\lambda}(p-q_1-k_2-k_1'-(-q_2+k_3'))^\eta + g^{\lambda\eta}(-q_2+k_3'-(k_2+k_2'))^\theta + g^{\eta\theta}(k_2+k_2'-(p-k_2-k_1'))^\lambda]
\end{aligned} \tag{163}$$

with

$$\begin{aligned} A_{c2} &= x_2 x_2'(r^2-1)m_B^2, & B_{c2} &= (r^2-1)x_3'(y-1)m_B^2 \\ C_{c2} &= (r^2-1)(x_2+y-1)(x_1'+1)m_B^2, & D_{c2} &= m_c^2+((r^2-1)x_1'+1)(x_3-y)m_B^2, \end{aligned} \tag{164}$$

and the color factor

$$C_N^{c2} = \varepsilon^{abc}\varepsilon^{a'b'c'} f_{jil}\left(C_1(T^jT^i)_{ac'}(T^l)_{bb'}\delta_{ca'} + C_2(T^i)_{cc'}(T^l)_{bb'}(T^j)_{aa'}\right) = 6iC_1. \tag{165}$$

Appendix C: The maximal one of $t_{1,2}$

The maximal one of the hard scales t_1^i and t_2^i for diagrams (a1)∼(a36) in Fig. 1, (b1)∼(b8) in Fig. 2 and (c1), (c2) in Fig. 3 are given below in the table.

$t_1^{a1} = max\{\sqrt{|A_{a1}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a1} = max\{\sqrt{|B_{a1}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a2} = max\{\sqrt{|A_{a2}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a2} = max\{\sqrt{|B_{a2}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a3} = max\{\sqrt{|A_{a3}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a3} = max\{\sqrt{|B_{a3}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a4} = max\{\sqrt{|A_{a4}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a4} = max\{\sqrt{|B_{a4}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a5} = max\{\sqrt{|A_{a5}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a5} = max\{\sqrt{|B_{a5}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a6} = max\{\sqrt{|A_{a6}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a6} = max\{\sqrt{|B_{a6}|}, \frac{1}{|\mathbf{b_1'}|}, \omega, \omega'\}$
$t_1^{a7} = max\{\sqrt{|A_{a7}|}, \frac{1}{|\mathbf{b_2}+\mathbf{b_1'}|}, \omega, \omega'\}$, $t_2^{a7} = max\{\sqrt{|B_{a7}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a8} = max\{\sqrt{|A_{a8}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a8} = max\{\sqrt{|B_{a8}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a9} = max\{\sqrt{|A_{a9}|}, \frac{1}{|\mathbf{b_1}|}, \omega, \omega'\}$, $t_2^{a9} = max\{\sqrt{|B_{a9}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a10} = max\{\sqrt{|A_{a10}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a10} = max\{\sqrt{|B_{a10}|}, \frac{1}{|\mathbf{b_3}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a11} = max\{\sqrt{|A_{a11}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a11} = max\{\sqrt{|B_{a11}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_2'}-\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a12} = max\{\sqrt{|A_{a12}|}, \frac{1}{|\mathbf{b_2}+\mathbf{b_3'}-\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a12} = max\{\sqrt{|B_{a12}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a13} = max\{\sqrt{|A_{a13}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a13} = max\{\sqrt{|B_{a13}|}, \frac{1}{|\mathbf{b_3}-\mathbf{b_1'}|}, \omega, \omega'\}$
$t_1^{a14} = max\{\sqrt{|A_{a14}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b_1'}|}, \omega, \omega'\}$, $t_2^{a14} = max\{\sqrt{|B_{a14}|}, \frac{1}{|\mathbf{b_3}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a15} = max\{\sqrt{|A_{a15}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a15} = max\{\sqrt{|B_{a15}|}, \frac{1}{|\mathbf{b_3}-\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a16} = max\{\sqrt{|A_{a16}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a16} = max\{\sqrt{|B_{a16}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_2'}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a17} = max\{\sqrt{|A_{a17}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a17} = max\{\sqrt{|B_{a17}|}, \frac{1}{|\mathbf{b_3}-\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a18} = max\{\sqrt{|A_{a18}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a18} = max\{\sqrt{|B_{a18}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a19} = max\{\sqrt{|A_{a19}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a19} = max\{\sqrt{|B_{a19}|}, \frac{1}{|\mathbf{b_3}-\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a20} = max\{\sqrt{|A_{a20}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a20} = max\{\sqrt{|B_{a20}|}, \frac{1}{|\mathbf{b_1}-\mathbf{b_1'}|}, \omega, \omega'\}$
$t_1^{a21} = max\{\sqrt{|A_{a21}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a21} = max\{\sqrt{|B_{a21}|}, \frac{1}{|\mathbf{b_1}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a22} = max\{\sqrt{|A_{a22}|}, \frac{1}{|\mathbf{b_2'}-\mathbf{b_3'}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a22} = max\{\sqrt{|B_{a22}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a23} = max\{\sqrt{|A_{a23}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a23} = max\{\sqrt{|B_{a23}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a24} = max\{\sqrt{|A_{a24}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a24} = max\{\sqrt{|B_{a24}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a25} = max\{\sqrt{|A_{a25}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a25} = max\{\sqrt{|B_{a25}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a26} = max\{\sqrt{|A_{a26}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a26} = max\{\sqrt{|B_{a26}|}, \frac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a27} = max\{\sqrt{|A_{a27}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a27} = max\{\sqrt{|B_{a27}|}, \frac{1}{|\mathbf{b_1'}|}, \omega, \omega'\}$
$t_1^{a28} = max\{\sqrt{|A_{a28}|}, \frac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$, $t_2^{a28} = max\{\sqrt{|B_{a28}|}, \frac{1}{|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}-\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a29} = max\{\sqrt{|A_{a29}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a29} = max\{\sqrt{|B_{a29}|}, \frac{1}{|\mathbf{b_1'}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a30} = max\{\sqrt{|A_{a30}|}, \frac{1}{|\mathbf{b_2}|}, \omega, \omega'\}$, $t_2^{a30} = max\{\sqrt{|B_{a30}|}, \frac{1}{|\mathbf{b_1'}+\mathbf{b_q}|}, \omega, \omega'\}$
$t_1^{a31} = max\{\sqrt{|A_{a31}|}, \frac{1}{|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_3'}+\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a31} = max\{\sqrt{|B_{a31}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$
$t_1^{a32} = max\{\sqrt{|A_{a32}|}, \frac{1}{|\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_3'}-\mathbf{b_q}|}, \omega, \omega'\}$, $t_2^{a32} = max\{\sqrt{|B_{a32}|}, \frac{1}{|\mathbf{b_3'}|}, \omega, \omega'\}$

$$t_1^{a33} = max\{\sqrt{|A_{a33}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{a33} = max\{\sqrt{|B_{a33}|}, \tfrac{1}{|\mathbf{b_3}-\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{a34} = max\{\sqrt{|A_{a34}|}, \tfrac{1}{|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{a34} = max\{\sqrt{|B_{a34}|}, \tfrac{1}{|\mathbf{b_1}-\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{a35} = max\{\sqrt{|A_{a35}|}, \tfrac{1}{|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_3'}|}, \omega, \omega'\}, \qquad t_2^{a35} = max\{\sqrt{|B_{a35}|}, \tfrac{1}{|\mathbf{b_3'}-\mathbf{b_q}|}, \omega, \omega'\}$$

$$t_1^{a36} = max\{\sqrt{|A_{a36}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{a36} = max\{\sqrt{|B_{a36}|}, \tfrac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}$$

$$t_1^{b1} = max\{\sqrt{|A_{b1}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{b1} = max\{\sqrt{|B_{b1}|}, \tfrac{1}{|\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{b2} = max\{\sqrt{|A_{b2}|}, \tfrac{1}{|\mathbf{b_2}+\mathbf{b_1'}|}, \omega, \omega'\}, \qquad t_2^{b2} = max\{\sqrt{|B_{b2}|}, \tfrac{1}{|\mathbf{b_q}|}, \omega, \omega'\}$$

$$t_1^{b3} = max\{\sqrt{|A_{b3}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{b3} = max\{\sqrt{|B_{b3}|}, \tfrac{1}{|\mathbf{b_3}-\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{b4} = max\{\sqrt{|A_{b4}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b_1'}|}, \omega, \omega'\}, \qquad t_2^{b4} = max\{\sqrt{|B_{b4}|}, \tfrac{1}{|\mathbf{b_3}+\mathbf{b_q}|}, \omega, \omega'\}$$

$$t_1^{b5} = max\{\sqrt{|A_{b5}|}, \tfrac{1}{|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{b5} = max\{\sqrt{|B_{b5}|}, \tfrac{1}{|\mathbf{b_1}-\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{b6} = max\{\sqrt{|A_{b6}|}, \tfrac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}, \qquad t_2^{b6} = max\{\sqrt{|B_{b6}|}, \tfrac{1}{|\mathbf{b_1}+\mathbf{b_q}|}, \omega, \omega'\}$$

$$t_1^{b7} = max\{\sqrt{|A_{b7}|}, \tfrac{1}{|\mathbf{b_2}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{b7} = max\{\sqrt{|B_{b7}|}, \tfrac{1}{|\mathbf{b_1'}|}, \omega, \omega'\}$$

$$t_1^{b8} = max\{\sqrt{|A_{b8}|}, \tfrac{1}{|\mathbf{b_2'}|}, \omega, \omega'\}, \qquad t_2^{b8} = max\{\sqrt{|B_{b8}|}, \tfrac{1}{|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}-\mathbf{b_q}|}, \omega, \omega'\}$$

$$t_1^{c1} = max\{\sqrt{|A_{c1}|}, \tfrac{2}{|\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}|}, \omega, \omega'\}, \qquad t_2^{c1} = max\{\sqrt{|B_{c1}|}, \tfrac{2}{|\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_2'}|}, \omega, \omega'\},$$

$$t_3^{c1} = max\{\sqrt{|C_{c1}|}, \tfrac{2}{|\mathbf{b_2}-\mathbf{b_1'}-2\mathbf{b_q}-\mathbf{b_2'}|}, \omega, \omega'\}$$

$$t_1^{c2} = max\{\sqrt{|A_{c2}|}, \tfrac{2}{|2\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_q}|}, \omega, \omega'\}, \qquad t_2^{c2} = max\{\sqrt{|B_{c2}|}, \tfrac{2}{|2\mathbf{b_3}-\mathbf{b_1'}+\mathbf{b_q}|}, \omega, \omega'\},$$

$$t_3^{c2} = max\{\sqrt{|C_{c2}|}, \tfrac{2}{|\mathbf{b_1'}+\mathbf{b_q}|}, \omega, \omega'\}$$

Appendix D: Expressions of Ω^i

The expression of Ω^i for diagrams (a1)∼(a36) in Fig. 1, (b1)∼(b8) in Fig. 2 and (c1), (c2) in Fig. 3 are presented below where K_0, K_1, N_0, N_1, J_0 and J_1 are Bessel functions and $\theta(x)$ is θ-function.

$$\begin{aligned}&\Omega^{a1} = \\ &\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)} \frac{\sqrt{X_2}}{\sqrt{|Z_2|}} \left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\ &\left[2\pi K_0(\sqrt{A_{a1}}|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|)\theta(A_{a1}) + \pi^2[-N_0(\sqrt{|A_{a1}|}|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a1}|}|\mathbf{b_2}-\mathbf{b_3'}+\mathbf{b_q}|)]\theta(-A_{a1})\right]\end{aligned} \tag{166}$$

with

$$\begin{aligned} Z_2 &= B_{a1}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[C_{a1}(1-z_1) + D_{a1}z_1] \\ X_2 &= [\mathbf{b_3'} - z_1(\mathbf{b_3'}-\mathbf{b_q})]^2 + \frac{z_1(1-z_1)}{z_2}(\mathbf{b_3'}-\mathbf{b_q})^2. \end{aligned} \tag{167}$$

$$\begin{aligned}&\Omega^{a2} = \\ &\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)} \frac{\sqrt{X_2}}{\sqrt{|Z_2|}} \left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\ &\left[2\pi K_0(\sqrt{B_{a2}}|\mathbf{b_q}|)\theta(B_{a2}) + \pi^2[-N_0(\sqrt{|B_{a2}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a2}|}|\mathbf{b_q}|)]\theta(-B_{a2})\right]\end{aligned} \tag{168}$$

with

$$\begin{aligned} Z_2 &= C_{a2}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{a2}(1-z_1) + D_{a2}z_1] \\ X_2 &= [\mathbf{b_3'} - \mathbf{b_q} - z_1\mathbf{b_2}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b_2}^2. \end{aligned} \tag{169}$$

$$
\begin{aligned}
&\Omega^{a3} = \\
&\left[2\pi K_0(\sqrt{A_{a3}}|\mathbf{b_2}|)\theta(A_{a3}) + \pi^2\left(-N_0(\sqrt{|A_{a3}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a3}|}|\mathbf{b_2}|)\right)\theta(-A_{a3})\right] \\
&\left[2\pi K_0(\sqrt{B_{a3}}|\mathbf{b_q}|)\theta(B_{a3}) + \pi^2\left(-N_0(\sqrt{|B_{a3}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a3}|}|\mathbf{b_q}|)\right)\theta(-B_{a3})\right] \\
&\left[2\pi K_0(\sqrt{C_{a3}}|\mathbf{b_3'} - \mathbf{b_q}|)\theta(C_{a3}) + \pi^2\left(-N_0(\sqrt{|C_{a3}|}|\mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a3}|}|\mathbf{b_3'} - \mathbf{b_q}|)\right)\theta(-C_{a3})\right] \\
&\left[2\pi K_0(\sqrt{D_{a3}}|\mathbf{b_3}|)\theta(D_{a3}) + \pi^2\left(-N_0(\sqrt{|D_{a3}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a3}|}|\mathbf{b_3}|)\right)\theta(-D_{a3})\right]
\end{aligned}
\tag{170}
$$

$$
\begin{aligned}
&\Omega^{a4} = \\
&\left[2\pi K_0(\sqrt{A_{a4}}|\mathbf{b_2}|)\theta(A_{a4}) + \pi^2\left(-N_0(\sqrt{|A_{a4}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a4}|}|\mathbf{b_2}|)\right)\theta(-A_{a4})\right] \\
&\left[2\pi K_0(\sqrt{B_{a4}}|\mathbf{b_q}|)\theta(B_{a4}) + \pi^2\left(-N_0(\sqrt{|B_{a4}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a4}|}|\mathbf{b_q}|)\right)\theta(-B_{a4})\right] \\
&\left[2\pi K_0(\sqrt{C_{a4}}|\mathbf{b_1}|)\theta(C_{a4}) + \pi^2\left(-N_0(\sqrt{|C_{a4}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|C_{a4}|}|\mathbf{b_1}|)\right)\theta(-C_{a4})\right] \\
&\left[2\pi K_0(\sqrt{D_{a4}}|\mathbf{b_3'} - \mathbf{b_q}|)\theta(D_{a4}) + \pi^2\left(-N_0(\sqrt{|D_{a4}|}|\mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|D_{a4}|}|\mathbf{b_3'} - \mathbf{b_q}|)\right)\theta(-D_{a4})\right]
\end{aligned}
\tag{171}
$$

$$
\begin{aligned}
&\Omega^{a5} = \\
&\left[2\pi K_0(\sqrt{A_{a5}}|\mathbf{b_2'}|)\theta(A_{a5}) + \pi^2\left(-N_0(\sqrt{|A_{a5}|}|\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{a5}|}|\mathbf{b_2'}|)\right)\theta(-A_{a5})\right] \\
&\left[2\pi K_0(\sqrt{B_{a5}}|\mathbf{b_q}|)\theta(B_{a5}) + \pi^2\left(-N_0(\sqrt{|B_{a5}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a5}|}|\mathbf{b_q}|)\right)\theta(-B_{a5})\right] \\
&\left[2\pi K_0(\sqrt{C_{a5}}|\mathbf{b_2} - \mathbf{b_2'}|)\theta(C_{a5}) + \pi^2\left(-N_0(\sqrt{|C_{a5}|}|\mathbf{b_2} - \mathbf{b_2'}|) + iJ_0(\sqrt{|C_{a5}|}|\mathbf{b_2} - \mathbf{b_2'}|)\right)\theta(-C_{a5})\right] \\
&[2\pi K_0(\sqrt{D_{a5}}|\mathbf{b_2} - \mathbf{b_2'} + \mathbf{b_3'} - \mathbf{b_q}|)\theta(D_{a5}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{a5}|}|\mathbf{b_2} - \mathbf{b_2'} + \mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|D_{a5}|}|\mathbf{b_2} - \mathbf{b_2'} + \mathbf{b_3'} - \mathbf{b_q}|)\right)\theta(-D_{a5})]
\end{aligned}
\tag{172}
$$

$$
\begin{aligned}
&\Omega^{a6} = \\
&\int_0^1 \frac{dz_1 dz_2 dz_3}{z_1(1-z_1)z_2(1-z_2)}\frac{\sqrt{X_3}}{\sqrt{|Z_3|}}\left[\pi^3 K_1(\sqrt{X_3 Z_3})\theta(Z_3) + \frac{\pi^4}{2}[N_1(\sqrt{X_3|Z_3|}) - iJ_1(\sqrt{X_3|Z_3|})]\theta(-Z_3)\right]
\end{aligned}
\tag{173}
$$

with

$$
\begin{aligned}
Z_3 &= B_{a6}(1-z_3) + \frac{z_3}{z_2(1-z_2)}\left[C_{a6}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{a6}(1-z_1) + D_{a6}z_1]\right] \\
X_3 &= [-\mathbf{b_1'} + z_2(\mathbf{b_1'} + \mathbf{b_q}) - (\mathbf{b_2} - \mathbf{b_q})z_2(1-z_1)]^2 + \frac{z_2(1-z_2)}{z_3}[-\mathbf{b_1'} - \mathbf{b_q} - (\mathbf{b_2} - \mathbf{b_q})z_1]^2.
\end{aligned}
\tag{174}
$$

$$
\begin{aligned}
&\Omega^{a7} = \\
&\int_0^1 dz\frac{|\mathbf{b_1'} + \mathbf{b_q}|}{\sqrt{|Z_1|}}\left[\pi K_1(\sqrt{Z_1}|\mathbf{b_1'} + \mathbf{b_q}|)\theta(Z_1) + \frac{\pi^2}{2}[N_1(\sqrt{|Z_1|}|\mathbf{b_1'} + \mathbf{b_q}|) - iJ_1(\sqrt{|Z_1|}|\mathbf{b_1'} + \mathbf{b_q}|)]\theta(-Z_1)\right] \\
&\left[2\pi K_0(\sqrt{A_{a7}}|\mathbf{b_2} + \mathbf{b_1'}|)\theta(A_{a7}) + \pi^2\left(-N_0(\sqrt{|A_{a7}|}|\mathbf{b_2} + \mathbf{b_1'}|) + iJ_0(\sqrt{|A_{a7}|}|\mathbf{b_2} + \mathbf{b_1'}|)\right)\theta(-A_{a7})\right] \\
&\left[2\pi K_0(\sqrt{B_{a7}}|\mathbf{b_q}|)\theta(B_{a7}) + \pi^2\left(-N_0(\sqrt{|B_{a7}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a7}|}|\mathbf{b_q}|)\right)\theta(-B_{a7})\right]
\end{aligned}
\tag{175}
$$

with

$$Z_1 = C_{a7}z + D_{a7}(1-z). \tag{176}$$

$$\begin{aligned}
&\Omega^{a8} = \\
&\left[2\pi K_0(\sqrt{A_{a8}}|\mathbf{b_2} - \mathbf{b_3'} + \mathbf{b_q}|)\theta(A_{a8}) + \pi^2\left(-N_0(\sqrt{|A_{a8}|}|\mathbf{b_2} - \mathbf{b_3'} + \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a8}|}|\mathbf{b_2} - \mathbf{b_3'} + \mathbf{b_q}|)\right)\theta(-A_{a8})\right] \\
&\left[2\pi K_0(\sqrt{B_{a8}}|\mathbf{b_1} - \mathbf{b_3'}|)\theta(B_{a8}) + \pi^2\left(-N_0(\sqrt{|B_{a8}|}|\mathbf{b_1} - \mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a8}|}|\mathbf{b_1} - \mathbf{b_3'}|)\right)\theta(-B_{a8})\right] \\
&\left[2\pi K_0(\sqrt{C_{a8}}|\mathbf{b_3'} - \mathbf{b_q}|)\theta(C_{a8}) + \pi^2\left(-N_0(\sqrt{|C_{a8}|}|\mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a8}|}|\mathbf{b_3'} - \mathbf{b_q}|)\right)\theta(-C_{a8})\right] \\
&\left[2\pi K_0(\sqrt{D_{a8}}|\mathbf{b_1}|)\theta(D_{a8}) + \pi^2\left(-N_0(\sqrt{|D_{a8}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a8}|}|\mathbf{b_1}|)\right)\theta(-D_{a8})\right]
\end{aligned} \tag{177}$$

$$\begin{aligned}
&\Omega^{a9} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)}\frac{\sqrt{X_2}}{\sqrt{|Z_2|}}\left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\left[2\pi K_0(\sqrt{B_{a9}}|\mathbf{b_1} - \mathbf{b_3'}|)\theta(B_{a9}) + \pi^2[-N_0(\sqrt{|B_{a9}|}|\mathbf{b_1} - \mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a9}|}|\mathbf{b_1} - \mathbf{b_3'}|)]\theta(-B_{a9})\right]
\end{aligned} \tag{178}$$

with

$$\begin{aligned}
Z_2 &= C_{a9}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{a9}(1-z_1) + D_{a9}z_1] \\
X_2 &= [\mathbf{b_1} - \mathbf{b_3'} + \mathbf{b_q} - z_1\mathbf{b_1}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b_1}^2.
\end{aligned} \tag{179}$$

$$\begin{aligned}
&\Omega^{a10} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)}\frac{\sqrt{X_2}}{\sqrt{|Z_2|}}\left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\left[2\pi K_0(\sqrt{A_{a10}}|\mathbf{b_2}|)\theta(A_{a10}) + \pi^2[-N_0(\sqrt{|A_{a10}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a10}|}|\mathbf{b_2}|)]\theta(-A_{a10})\right]
\end{aligned} \tag{180}$$

with

$$\begin{aligned}
Z_2 &= B_{a10}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[C_{a10}(1-z_1) + D_{a10}z_1] \\
X_2 &= [\mathbf{b_3} + \mathbf{b_q} - z_1\mathbf{b_3}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b_3}^2.
\end{aligned} \tag{181}$$

$$\begin{aligned}
&\Omega^{a11} = \\
&\left[2\pi K_0(\sqrt{A_{a11}}|\mathbf{b_2}|)\theta(A_{a11}) + \pi^2\left(-N_0(\sqrt{|A_{a11}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a11}|}|\mathbf{b_2}|)\right)\theta(-A_{a11})\right] \\
&[2\pi K_0(\sqrt{B_{a11}}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'} - \mathbf{b_q}|)\theta(B_{a11}) + \\
&\pi^2\left(-N_0(\sqrt{|B_{a11}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'} - \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a11}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'} - \mathbf{b_q}|)\right)\theta(-B_{a11})] \\
&\left[2\pi K_0(\sqrt{C_{a11}}|\mathbf{b_2} - \mathbf{b_2'}|)\theta(C_{a11}) + \pi^2\left(-N_0(\sqrt{|C_{a11}|}|\mathbf{b_2} - \mathbf{b_2'}|) + iJ_0(\sqrt{|C_{a11}|}|\mathbf{b_2} - \mathbf{b_2'}|)\right)\theta(-C_{a11})\right] \\
&\left[2\pi K_0(\sqrt{D_{a11}}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'}|)\theta(D_{a11}) + \pi^2\left(-N_0(\sqrt{|D_{a11}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'}|) + iJ_0(\sqrt{|D_{a11}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_2'}|)\right)\theta(-D_{a11})\right]
\end{aligned} \tag{182}$$

$$\begin{aligned}
&\Omega^{a12} = \\
&\Big[2\pi K_0(\sqrt{A_{a12}}|\mathbf{b_2} + \mathbf{b'_3} - \mathbf{b_q}|)\theta(A_{a12}) + \pi^2\left(-N_0(\sqrt{|A_{a12}|}|\mathbf{b_2} + \mathbf{b'_3} - \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a12}|}|\mathbf{b_2} + \mathbf{b'_3} - \mathbf{b_q}|)\right)\theta(-A_{a12})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a12}}|\mathbf{b_1} - \mathbf{b_q}|)\theta(B_{a12}) + \pi^2\left(-N_0(\sqrt{|B_{a12}|}|\mathbf{b_1} - \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a12}|}|\mathbf{b_1} - \mathbf{b_q}|)\right)\theta(-B_{a12})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a12}}|\mathbf{b'_3} - \mathbf{b_q}|)\theta(C_{a12}) + \pi^2\left(-N_0(\sqrt{|C_{a12}|}|\mathbf{b'_3} - \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a12}|}|\mathbf{b'_3} - \mathbf{b_q}|)\right)\theta(-C_{a12})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{a12}}|\mathbf{b_1}|)\theta(D_{a12}) + \pi^2\left(-N_0(\sqrt{|D_{a12}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a12}|}|\mathbf{b_1}|)\right)\theta(-D_{a12})\Big]
\end{aligned} \tag{183}$$

$$\begin{aligned}
&\Omega^{a13} = \\
&\Big[2\pi K_0(\sqrt{A_{a13}}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_q}|)\theta(A_{a13}) + \pi^2\left(-N_0(\sqrt{|A_{a13}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a13}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_q}|)\right)\theta(-A_{a13})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a13}}|\mathbf{b_3} - \mathbf{b'_1}|)\theta(B_{a13}) + \pi^2\left(-N_0(\sqrt{|B_{a13}|}|\mathbf{b_3} - \mathbf{b'_1}|) + iJ_0(\sqrt{|B_{a13}|}|\mathbf{b_3} - \mathbf{b'_1}|)\right)\theta(-B_{a13})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a13}}|\mathbf{b'_1} + \mathbf{b_q}|)\theta(C_{a13}) + \pi^2\left(-N_0(\sqrt{|C_{a13}|}|\mathbf{b'_1} + \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a13}|}|\mathbf{b'_1} + \mathbf{b_q}|)\right)\theta(-C_{a13})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{a13}}|\mathbf{b_3}|)\theta(D_{a13}) + \pi^2\left(-N_0(\sqrt{|D_{a13}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a13}|}|\mathbf{b_3}|)\right)\theta(-D_{a13})\Big]
\end{aligned} \tag{184}$$

$$\begin{aligned}
&\Omega^{a14} = \\
&\Big[2\pi K_0(\sqrt{A_{a14}}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b'_1}|)\theta(A_{a14}) + \pi^2\left(-N_0(\sqrt{|A_{a14}|}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b'_1}|) + iJ_0(\sqrt{|A_{a14}|}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b'_1}|)\right)\theta(-A_{a14})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a14}}|\mathbf{b_3} + \mathbf{b_q}|)\theta(B_{a14}) + \pi^2\left(-N_0(\sqrt{|B_{a14}|}|\mathbf{b_3} + \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a14}|}|\mathbf{b_3} + \mathbf{b_q}|)\right)\theta(-B_{a14})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a14}}|\mathbf{b'_1} + \mathbf{b_q}|)\theta(C_{a14}) + \pi^2\left(-N_0(\sqrt{|C_{a14}|}|\mathbf{b'_1} + \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a14}|}|\mathbf{b'_1} + \mathbf{b_q}|)\right)\theta(-C_{a14})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{a14}}|\mathbf{b_3}|)\theta(D_{a14}) + \pi^2\left(-N_0(\sqrt{|D_{a14}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a14}|}|\mathbf{b_3}|)\right)\theta(-D_{a14})\Big]
\end{aligned} \tag{185}$$

$$\begin{aligned}
&\Omega^{a15} = \\
&\Big[2\pi K_0(\sqrt{A_{a15}}|\mathbf{b_2} - \mathbf{b'_3} + \mathbf{b_q}|)\theta(A_{a15}) + \pi^2\left(-N_0(\sqrt{|A_{a15}|}|\mathbf{b_2} - \mathbf{b'_3} + \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a15}|}|\mathbf{b_2} - \mathbf{b'_3} + \mathbf{b_q}|)\right)\theta(-A_{a15})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a15}}|\mathbf{b_3} - \mathbf{b'_3}|)\theta(B_{a15}) + \pi^2\left(-N_0(\sqrt{|B_{a15}|}|\mathbf{b_3} - \mathbf{b'_3}|) + iJ_0(\sqrt{|B_{a15}|}|\mathbf{b_3} - \mathbf{b'_3}|)\right)\theta(-B_{a15})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a15}}|\mathbf{b_3}|)\theta(C_{a15}) + \pi^2\left(-N_0(\sqrt{|C_{a15}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|C_{a15}|}|\mathbf{b_3}|)\right)\theta(-C_{a15})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{a15}}|\mathbf{b'_3} - \mathbf{b_q}|)\theta(D_{a15}) + \pi^2\left(-N_0(\sqrt{|D_{a15}|}|\mathbf{b'_3} - \mathbf{b_q}|) + iJ_0(\sqrt{|D_{a15}|}|\mathbf{b'_3} - \mathbf{b_q}|)\right)\theta(-D_{a15})\Big]
\end{aligned} \tag{186}$$

$$
\begin{aligned}
&\Omega^{a16} = \\
&\left[2\pi K_0(\sqrt{A_{a16}}|\mathbf{b_2}|)\theta(A_{a16}) + \pi^2\left(-N_0(\sqrt{|A_{a16}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a16}|}|\mathbf{b_2}|)\right)\theta(-A_{a16})\right] \\
&[2\pi K_0(\sqrt{B_{a16}}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'} + \mathbf{b_q}|)\theta(B_{a16}) + \\
&\pi^2\left(-N_0(\sqrt{|B_{a16}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'} + \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a16}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'} + \mathbf{b_q}|)\right)\theta(-B_{a16})] \\
&\left[2\pi K_0(\sqrt{C_{a16}}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'}|)\theta(C_{a16}) + \pi^2\left(-N_0(\sqrt{|C_{a16}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'}|) + iJ_0(\sqrt{|C_{a16}|}|\mathbf{b_2} - \mathbf{b_3} - \mathbf{b_2'}|)\right)\theta(-C_{a16})\right] \\
&\left[2\pi K_0(\sqrt{D_{a16}}|\mathbf{b_2} - \mathbf{b_2'}|)\theta(D_{a16}) + \pi^2\left(-N_0(\sqrt{|D_{a16}|}|\mathbf{b_2} - \mathbf{b_2'}|) + iJ_0(\sqrt{|D_{a16}|}|\mathbf{b_2} - \mathbf{b_2'}|)\right)\theta(-D_{a16})\right]
\end{aligned}
\tag{187}
$$

$$
\begin{aligned}
&\Omega^{a17} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)} \frac{\sqrt{X_2}}{\sqrt{|Z_2|}} \left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\left[2\pi K_0(\sqrt{B_{a17}}|\mathbf{b_3} - \mathbf{b_q}|)\theta(B_{a17}) + \pi^2[-N_0(\sqrt{|B_{a17}|}|\mathbf{b_3} - \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a17}|}|\mathbf{b_3} - \mathbf{b_q}|)]\theta(-B_{a17})\right]
\end{aligned}
\tag{188}
$$

with

$$
\begin{aligned}
Z_2 &= C_{a17}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{a17}(1-z_1) + D_{a17}z_1] \\
X_2 &= [\mathbf{b_3} - z_1\mathbf{b_2}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b_2}^2.
\end{aligned}
\tag{189}
$$

$$
\begin{aligned}
&\Omega^{a18} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)} \frac{\sqrt{X_2}}{\sqrt{|Z_2|}} \left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\left[2\pi K_0(\sqrt{A_{a18}}|\mathbf{b_2}|)\theta(A_{a18}) + \pi^2[-N_0(\sqrt{|A_{a18}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a18}|}|\mathbf{b_2}|)]\theta(-A_{a18})\right]
\end{aligned}
\tag{190}
$$

with

$$
\begin{aligned}
Z_2 &= B_{a18}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[C_{a18}(1-z_1) + D_{a18}z_1] \\
X_2 &= [-\mathbf{b_q} - z_1\mathbf{b_1}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b_1}^2.
\end{aligned}
\tag{191}
$$

$$
\begin{aligned}
&\Omega^{a19} = \\
&\left[2\pi K_0(\sqrt{A_{a19}}|\mathbf{b_2'}|)\theta(A_{a19}) + \pi^2\left(-N_0(\sqrt{|A_{a19}|}|\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{a19}|}|\mathbf{b_2'}|)\right)\theta(-A_{a19})\right] \\
&\left[2\pi K_0(\sqrt{B_{a19}}|-\mathbf{b_3} + \mathbf{b_q}|)\theta(B_{a19}) + \pi^2\left(-N_0(\sqrt{|B_{a19}|}|-\mathbf{b_3} + \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a19}|}|-\mathbf{b_3} + \mathbf{b_q}|)\right)\theta(-B_{a19})\right] \\
&\left[2\pi K_0(\sqrt{C_{a19}}|\mathbf{b_3}|)\theta(C_{a19}) + \pi^2\left(-N_0(\sqrt{|C_{a19}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|C_{a19}|}|\mathbf{b_3}|)\right)\theta(-C_{a19})\right] \\
&\left[2\pi K_0(\sqrt{D_{a19}}|\mathbf{b_2} - \mathbf{b_2'}|)\theta(D_{a19}) + \pi^2\left(-N_0(\sqrt{|D_{a19}|}|\mathbf{b_2} - \mathbf{b_2'}|) + iJ_0(\sqrt{|D_{a19}|}|\mathbf{b_2} - \mathbf{b_2'}|)\right)\theta(-D_{a19})\right]
\end{aligned}
\tag{192}
$$

$$\begin{aligned}
&\Omega^{a20} = \\
&\left[2\pi K_0(\sqrt{A_{a20}}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|)\theta(A_{a20}) + \pi^2\left(-N_0(\sqrt{|A_{a20}|}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a20}|}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|)\right)\theta(-A_{a20})\right] \\
&\left[2\pi K_0(\sqrt{B_{a20}}|\mathbf{b_1}-\mathbf{b_1'}|)\theta(B_{a20}) + \pi^2\left(-N_0(\sqrt{|B_{a20}|}|\mathbf{b_1}-\mathbf{b_1'}|) + iJ_0(\sqrt{|B_{a20}|}|\mathbf{b_1}-\mathbf{b_1'}|)\right)\theta(-B_{a20})\right] \\
&\left[2\pi K_0(\sqrt{C_{a20}}|\mathbf{b_1'}+\mathbf{b_q}|)\theta(C_{a20}) + \pi^2\left(-N_0(\sqrt{|C_{a20}|}|\mathbf{b_1'}+\mathbf{b_q}|) + iJ_0(\sqrt{|C_{a20}|}|\mathbf{b_1'}+\mathbf{b_q}|)\right)\theta(-C_{a20})\right] \\
&\left[2\pi K_0(\sqrt{D_{a20}}|\mathbf{b_1}|)\theta(D_{a20}) + \pi^2\left(-N_0(\sqrt{|D_{a20}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a20}|}|\mathbf{b_1}|)\right)\theta(-D_{a20})\right]
\end{aligned}\tag{193}$$

$$\begin{aligned}
&\Omega^{a21} = \\
&\left[2\pi K_0(\sqrt{A_{a21}}|\mathbf{b_2'}|)\theta(A_{a21}) + \pi^2\left(-N_0(\sqrt{|A_{a21}|}|\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{a21}|}|\mathbf{b_2'}|)\right)\theta(-A_{a21})\right] \\
&\left[2\pi K_0(\sqrt{B_{a21}}|\mathbf{b_1}+\mathbf{b_q}|)\theta(B_{a21}) + \pi^2\left(-N_0(\sqrt{|B_{a21}|}|\mathbf{b_1}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a21}|}|\mathbf{b_1}+\mathbf{b_q}|)\right)\theta(-B_{a21})\right] \\
&\left[2\pi K_0(\sqrt{C_{a21}}|\mathbf{b_1'}+\mathbf{b_q}|)\theta(C_{a21}) + \pi^2\left(-N_0(\sqrt{|C_{a21}|}|\mathbf{b_1'}+\mathbf{b_q}|) + iJ_0(\sqrt{|C_{a21}|}|\mathbf{b_1'}+\mathbf{b_q}|)\right)\theta(-C_{a21})\right] \\
&\left[2\pi K_0(\sqrt{D_{a21}}|\mathbf{b_1}|)\theta(D_{a21}) + \pi^2\left(-N_0(\sqrt{|D_{a21}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a21}|}|\mathbf{b_1}|)\right)\theta(-D_{a21})\right]
\end{aligned}\tag{194}$$

$$\begin{aligned}
&\Omega^{a22} = \\
&\left[2\pi K_0(\sqrt{A_{a22}}|\mathbf{b_2'}-\mathbf{b_3'}+\mathbf{b_q}|)\theta(A_{a22}) + \pi^2\left(-N_0(\sqrt{|A_{a22}|}|\mathbf{b_2'}-\mathbf{b_3'}+\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a22}|}|\mathbf{b_2'}-\mathbf{b_3'}+\mathbf{b_q}|)\right)\theta(-A_{a22})\right] \\
&\left[2\pi K_0(\sqrt{B_{a22}}|\mathbf{b_3'}|)\theta(B_{a22}) + \pi^2\left(-N_0(\sqrt{|B_{a22}|}|\mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a22}|}|\mathbf{b_3'}|)\right)\theta(-B_{a22})\right] \\
&\left[2\pi K_0(\sqrt{C_{a22}}|\mathbf{b_2}-\mathbf{b_2'}|)\theta(C_{a22}) + \pi^2\left(-N_0(\sqrt{|C_{a22}|}|\mathbf{b_2}-\mathbf{b_2'}|) + iJ_0(\sqrt{|C_{a22}|}|\mathbf{b_2}-\mathbf{b_2'}|)\right)\theta(-C_{a22})\right] \\
&[2\pi K_0(\sqrt{D_{a22}}|\mathbf{b_2}-\mathbf{b_2'}+\mathbf{b_3'}-\mathbf{b_q}|)\theta(D_{a22}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{a22}|}|\mathbf{b_2}-\mathbf{b_2'}+\mathbf{b_3'}-\mathbf{b_q}|) + iJ_0(\sqrt{|D_{a22}|}|\mathbf{b_2}-\mathbf{b_2'}+\mathbf{b_3'}-\mathbf{b_q}|)\right)\theta(-D_{a22})]
\end{aligned}\tag{195}$$

$$\begin{aligned}
&\Omega^{a23} = \\
&\left[2\pi K_0(\sqrt{A_{a23}}|\mathbf{b_2}|)\theta(A_{a23}) + \pi^2\left(-N_0(\sqrt{|A_{a23}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a23}|}|\mathbf{b_2}|)\right)\theta(-A_{a23})\right] \\
&\left[2\pi K_0(\sqrt{B_{a23}}|\mathbf{b_3'}|)\theta(B_{a23}) + \pi^2\left(-N_0(\sqrt{|B_{a23}|}|\mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a23}|}|\mathbf{b_3'}|)\right)\theta(-B_{a23})\right] \\
&\left[2\pi K_0(\sqrt{C_{a23}}|\mathbf{b_3'}-\mathbf{b_q}|)\theta(C_{a23}) + \pi^2\left(-N_0(\sqrt{|C_{a23}|}|\mathbf{b_3'}-\mathbf{b_q}|) + iJ_0(\sqrt{|C_{a23}|}|\mathbf{b_3'}-\mathbf{b_q}|)\right)\theta(-C_{a23})\right] \\
&\left[2\pi K_0(\sqrt{D_{a23}}|\mathbf{b_3}|)\theta(D_{a23}) + \pi^2\left(-N_0(\sqrt{|D_{a23}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a23}|}|\mathbf{b_3}|)\right)\theta(-D_{a23})\right]
\end{aligned}\tag{196}$$

$$\begin{aligned}
&\Omega^{a24} = \\
&\Big[2\pi K_0(\sqrt{A_{a24}}|\mathbf{b_2}|)\theta(A_{a24}) + \pi^2\left(-N_0(\sqrt{|A_{a24}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a24}|}|\mathbf{b_2}|)\right)\theta(-A_{a24})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a24}}|\mathbf{b'_3}|)\theta(B_{a24}) + \pi^2\left(-N_0(\sqrt{|B_{a24}|}|\mathbf{b'_3}|) + iJ_0(\sqrt{|B_{a24}|}|\mathbf{b'_3}|)\right)\theta(-B_{a24})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a24}}|\mathbf{b'_3}-\mathbf{b_q}|)\theta(C_{a24}) + \pi^2\left(-N_0(\sqrt{|C_{a24}|}|\mathbf{b'_3}-\mathbf{b_q}|) + iJ_0(\sqrt{|C_{a24}|}|\mathbf{b'_3}-\mathbf{b_q}|)\right)\theta(-C_{a24})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{a24}}|\mathbf{b_1}|)\theta(D_{a24}) + \pi^2\left(-N_0(\sqrt{|D_{a24}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a24}|}|\mathbf{b_1}|)\right)\theta(-D_{a24})\Big]
\end{aligned} \tag{197}$$

$$\begin{aligned}
&\Omega^{a25} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)}\frac{\sqrt{X_2}}{\sqrt{|Z_2|}}\left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\Big[2\pi K_0(\sqrt{B_{a25}}|\mathbf{b'_3}|)\theta(B_{a25}) + \pi^2[-N_0(\sqrt{|B_{a25}|}|\mathbf{b'_3}|) + iJ_0(\sqrt{|B_{a25}|}|\mathbf{b'_3}|)]\theta(-B_{a25})\Big]
\end{aligned} \tag{198}$$

with

$$\begin{aligned}
Z_2 &= D_{a25}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{a25}(1-z_1) + C_{a25}z_1] \\
X_2 &= [\mathbf{b'_3} - \mathbf{b_q} - z_1\mathbf{b'_2}]^2 + \frac{z_1(1-z_1)}{z_2}\mathbf{b'_2}{}^2.
\end{aligned} \tag{199}$$

$$\begin{aligned}
&\Omega^{a26} = \\
&\int_0^1 \frac{dz_1 dz_2}{z_1(1-z_1)}\frac{\sqrt{X_2}}{\sqrt{|Z_2|}}\left[\pi^2 K_1(\sqrt{X_2 Z_2})\theta(Z_2) + \frac{\pi^3}{2}[N_1(\sqrt{X_2|Z_2|}) - iJ_1(\sqrt{X_2|Z_2|})]\theta(-Z_2)\right] \\
&\Big[2\pi K_0(\sqrt{A_{a26}}|\mathbf{b'_2}|)\theta(A_{a26}) + \pi^2[-N_0(\sqrt{|A_{a26}|}|\mathbf{b'_2}|) + iJ_0(\sqrt{|A_{a26}|}|\mathbf{b'_2}|)]\theta(-A_{a26})\Big]
\end{aligned} \tag{200}$$

with

$$\begin{aligned}
Z_2 &= B_{a26}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[D_{a26}(1-z_1) + C_{a26}z_1] \\
X_2 &= [\mathbf{b_q} - z_1(-\mathbf{b'_3} + \mathbf{b_q})]^2 + \frac{z_1(1-z_1)}{z_2}(\mathbf{b'_3} - \mathbf{b_q})^2.
\end{aligned} \tag{201}$$

$$\begin{aligned}
&\Omega^{a27} = \\
&\Big[2\pi K_0(\sqrt{A_{a27}}|\mathbf{b_2}-\mathbf{b_q}|)\theta(A_{a27}) + \pi^2\left(-N_0(\sqrt{|A_{a27}|}|\mathbf{b_2}-\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a27}|}|\mathbf{b_2}-\mathbf{b_q}|)\right)\theta(-A_{a27})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{a27}}|\mathbf{b'_1}|)\theta(B_{a27}) + \pi^2\left(-N_0(\sqrt{|B_{a27}|}|\mathbf{b'_1}|) + iJ_0(\sqrt{|B_{a27}|}|\mathbf{b'_1}|)\right)\theta(-B_{a27})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{a27}}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|)\theta(C_{a27}) + \pi^2\left(-N_0(\sqrt{|C_{a27}|}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|) + iJ_0(\sqrt{|C_{a27}|}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|)\right)\theta(-C_{a27})\Big] \\
&[2\pi K_0(\sqrt{D_{a27}}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|)\theta(D_{a27}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{a27}|}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|) + iJ_0(\sqrt{|D_{a27}|}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|)\right)\theta(-D_{a27})]
\end{aligned} \tag{202}$$

$$
\begin{aligned}
&\Omega^{a28} = \\
&\left[2\pi K_0(\sqrt{A_{a28}}|\mathbf{b_2'}|)\theta(A_{a28}) + \pi^2\left(-N_0(\sqrt{|A_{a28}|}|\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{a28}|}|\mathbf{b_2'}|)\right)\theta(-A_{a28})\right] \\
&[2\pi K_0(\sqrt{B_{a28}}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|)\theta(B_{a28}) + \\
&\pi^2\left(-N_0(\sqrt{|B_{a28}|}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a28}|}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|)\right)\theta(-B_{a28})] \\
&\left[2\pi K_0(\sqrt{C_{a28}}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|)\theta(C_{a28}) + \pi^2\left(-N_0(\sqrt{|C_{a28}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|) + iJ_0(\sqrt{|C_{a28}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|)\right)\theta(-C_{a28})\right] \\
&[2\pi K_0(\sqrt{D_{a28}}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|)\theta(D_{a28}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{a28}|}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|) + iJ_0(\sqrt{|D_{a28}|}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|)\right)\theta(-D_{a28})]
\end{aligned}
\tag{203}
$$

$$
\begin{aligned}
&\Omega^{a29} = \\
&\left[2\pi K_0(\sqrt{A_{a29}}|\mathbf{b_2}|)\theta(A_{a29}) + \pi^2\left(-N_0(\sqrt{|A_{a29}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a29}|}|\mathbf{b_2}|)\right)\theta(-A_{a29})\right] \\
&\left[2\pi K_0(\sqrt{B_{a29}}|\mathbf{b_1'}+\mathbf{b_q}|)\theta(B_{a29}) + \pi^2\left(-N_0(\sqrt{|B_{a29}|}|\mathbf{b_1'}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a29}|}|\mathbf{b_1'}+\mathbf{b_q}|)\right)\theta(-B_{a29})\right] \\
&\left[2\pi K_0(\sqrt{C_{a29}}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b_1'}|)\theta(C_{a29}) + \pi^2\left(-N_0(\sqrt{|C_{a29}|}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a29}|}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b_1'}|)\right)\theta(-C_{a29})\right] \\
&\left[2\pi K_0(\sqrt{D_{a29}}|\mathbf{b_3}|)\theta(D_{a29}) + \pi^2\left(-N_0(\sqrt{|D_{a29}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a29}|}|\mathbf{b_3}|)\right)\theta(-D_{a29})\right]
\end{aligned}
\tag{204}
$$

$$
\begin{aligned}
&\Omega^{a30} = \\
&\left[2\pi K_0(\sqrt{A_{a30}}|\mathbf{b_2}|)\theta(A_{a30}) + \pi^2\left(-N_0(\sqrt{|A_{a30}|}|\mathbf{b_2}|) + iJ_0(\sqrt{|A_{a30}|}|\mathbf{b_2}|)\right)\theta(-A_{a30})\right] \\
&\left[2\pi K_0(\sqrt{B_{a30}}|\mathbf{b_1'}+\mathbf{b_q}|)\theta(B_{a30}) + \pi^2\left(-N_0(\sqrt{|B_{a30}|}|\mathbf{b_1'}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{a30}|}|\mathbf{b_1'}+\mathbf{b_q}|)\right)\theta(-B_{a30})\right] \\
&[2\pi K_0(\sqrt{C_{a30}}|-\mathbf{b_1}+\mathbf{b_2}+\mathbf{b_1'}|)\theta(C_{a30}) + \\
&\pi^2\left(-N_0(\sqrt{|C_{a30}|}|-\mathbf{b_1}+\mathbf{b_2}+\mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a30}|}|-\mathbf{b_1}+\mathbf{b_2}+\mathbf{b_1'}|)\right)\theta(-C_{a30})] \\
&\left[2\pi K_0(\sqrt{D_{a30}}|\mathbf{b_1}|)\theta(D_{a30}) + \pi^2\left(-N_0(\sqrt{|D_{a30}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a30}|}|\mathbf{b_1}|)\right)\theta(-D_{a30})\right]
\end{aligned}
\tag{205}
$$

$$
\begin{aligned}
&\Omega^{a31} = \\
&[2\pi K_0(\sqrt{A_{a31}}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_3'}+\mathbf{b_q}|)\theta(A_{a31}) + \\
&\pi^2\left(-N_0(\sqrt{|A_{a31}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_3'}+\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a31}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_3'}+\mathbf{b_q}|)\right)\theta(-A_{a31})] \\
&\left[2\pi K_0(\sqrt{B_{a31}}|\mathbf{b_3'}|)\theta(B_{a31}) + \pi^2\left(-N_0(\sqrt{|B_{a31}|}|\mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a31}|}|\mathbf{b_3'}|)\right)\theta(-B_{a31})\right] \\
&\left[2\pi K_0(\sqrt{C_{a31}}|\mathbf{b_2}+\mathbf{b_1'}|)\theta(C_{a31}) + \pi^2\left(-N_0(\sqrt{|C_{a31}|}|\mathbf{b_2}+\mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a31}|}|\mathbf{b_2}+\mathbf{b_1'}|)\right)\theta(-C_{a31})\right] \\
&[2\pi K_0(\sqrt{D_{a31}}|-\mathbf{b_1'}+\mathbf{b_3'}-\mathbf{b_q}|)\theta(D_{a31}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{a31}|}|-\mathbf{b_1'}+\mathbf{b_3'}-\mathbf{b_q}|) + iJ_0(\sqrt{|D_{a31}|}|-\mathbf{b_1'}+\mathbf{b_3'}-\mathbf{b_q}|)\right)\theta(-D_{a31})]
\end{aligned}
\tag{206}
$$

$$
\begin{aligned}
&\Omega^{a32} = \\
&[2\pi K_0(\sqrt{A_{a32}}|\mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|)\theta(A_{a32}) + \\
&\pi^2 \left(-N_0(\sqrt{|A_{a32}|}|\mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a32}|}|\mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|)\right) \theta(-A_{a32})] \\
&\left[2\pi K_0(\sqrt{B_{a32}}|\mathbf{b_3'}|)\theta(B_{a32}) + \pi^2 \left(-N_0(\sqrt{|B_{a32}|}|\mathbf{b_3'}|) + iJ_0(\sqrt{|B_{a32}|}|\mathbf{b_3'}|)\right) \theta(-B_{a32})\right] \\
&\left[2\pi K_0(\sqrt{C_{a32}}|\mathbf{b_1'} - \mathbf{b_3'} + \mathbf{b_q}|)\theta(C_{a32}) + \pi^2 \left(-N_0(\sqrt{|C_{a32}|}|\mathbf{b_1'} - \mathbf{b_3'} + \mathbf{b_q}|) + iJ_0(\sqrt{|C_{a32}|}|\mathbf{b_1'} - \mathbf{b_3'} + \mathbf{b_q}|)\right) \theta(-C_{a32})\right] \\
&\left[2\pi K_0(\sqrt{D_{a32}}|\mathbf{b_2} + \mathbf{b_3'} - \mathbf{b_q}|)\theta(D_{a32}) + \pi^2 \left(-N_0(\sqrt{|D_{a32}|}|\mathbf{b_2} + \mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|D_{a32}|}|\mathbf{b_2} + \mathbf{b_3'} - \mathbf{b_q}|)\right) \theta(-D_{a32})\right]
\end{aligned}
\tag{207}
$$

$$
\begin{aligned}
&\Omega^{a33} = \\
&[2\pi K_0(\sqrt{A_{a33}}| - \mathbf{b_2} + \mathbf{b_3} + \mathbf{b_q}|)\theta(A_{a33}) + \\
&\pi^2 \left(-N_0(\sqrt{|A_{a33}|}| - \mathbf{b_2} + \mathbf{b_3} + \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a33}|}| - \mathbf{b_2} + \mathbf{b_3} + \mathbf{b_q}|)\right) \theta(-A_{a33})] \\
&\left[2\pi K_0(\sqrt{B_{a33}}|\mathbf{b_3} - \mathbf{b_1'}|)\theta(B_{a33}) + \pi^2 \left(-N_0(\sqrt{|B_{a33}|}|\mathbf{b_3} - \mathbf{b_1'}|) + iJ_0(\sqrt{|B_{a33}|}|\mathbf{b_3} - \mathbf{b_1'}|)\right) \theta(-B_{a33})\right] \\
&\left[2\pi K_0(\sqrt{C_{a33}}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b_1'}|)\theta(C_{a33}) + \pi^2 \left(-N_0(\sqrt{|C_{a33}|}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a33}|}|\mathbf{b_2} - \mathbf{b_3} + \mathbf{b_1'}|)\right) \theta(-C_{a33})\right] \\
&\left[2\pi K_0(\sqrt{D_{a33}}|\mathbf{b_3}|)\theta(D_{a33}) + \pi^2 \left(-N_0(\sqrt{|D_{a33}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{a33}|}|\mathbf{b_3}|)\right) \theta(-D_{a33})\right]
\end{aligned}
\tag{208}
$$

$$
\begin{aligned}
&\Omega^{a34} = \\
&[2\pi K_0(\sqrt{A_{a34}}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_q}|)\theta(A_{a34}) + \\
&\pi^2 \left(-N_0(\sqrt{|A_{a34}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_q}|) + iJ_0(\sqrt{|A_{a34}|}|\mathbf{b_1} - \mathbf{b_2} + \mathbf{b_q}|)\right) \theta(-A_{a34})] \\
&\left[2\pi K_0(\sqrt{B_{a34}}|\mathbf{b_1} - \mathbf{b_1'}|)\theta(B_{a34}) + \pi^2 \left(-N_0(\sqrt{|B_{a34}|}|\mathbf{b_1} - \mathbf{b_1'}|) + iJ_0(\sqrt{|B_{a34}|}|\mathbf{b_1} - \mathbf{b_1'}|)\right) \theta(-B_{a34})\right] \\
&[2\pi K_0(\sqrt{C_{a34}}| - \mathbf{b_1} + \mathbf{b_2} + \mathbf{b_1'}|)\theta(C_{a34}) + \\
&\pi^2 \left(-N_0(\sqrt{|C_{a34}|}| - \mathbf{b_1} + \mathbf{b_2} + \mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a34}|}| - \mathbf{b_1} + \mathbf{b_2} + \mathbf{b_1'}|)\right) \theta(-C_{a34})] \\
&\left[2\pi K_0(\sqrt{D_{a34}}|\mathbf{b_1}|)\theta(D_{a34}) + \pi^2 \left(-N_0(\sqrt{|D_{a34}|}|\mathbf{b_1}|) + iJ_0(\sqrt{|D_{a34}|}|\mathbf{b_1}|)\right) \theta(-D_{a34})\right]
\end{aligned}
\tag{209}
$$

$$
\begin{aligned}
&\Omega^{a35} = \\
&[2\pi K_0(\sqrt{A_{a35}}| - \mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'}|)\theta(A_{a35}) + \\
&\pi^2 \left(-N_0(\sqrt{|A_{a35}|}| - \mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'}|) + iJ_0(\sqrt{|A_{a35}|}| - \mathbf{b_2} - \mathbf{b_1'} + \mathbf{b_3'}|)\right) \theta(-A_{a35})] \\
&\left[2\pi K_0(\sqrt{B_{a35}}| - \mathbf{b_3'} + \mathbf{b_q}|)\theta(B_{a35}) + \pi^2 \left(-N_0(\sqrt{|B_{a35}|}| - \mathbf{b_3'} + \mathbf{b_q}|) + iJ_0(\sqrt{|B_{a35}|}| - \mathbf{b_3'} + \mathbf{b_q}|)\right) \theta(-B_{a35})\right] \\
&\left[2\pi K_0(\sqrt{C_{a35}}|\mathbf{b_2} + \mathbf{b_1'}|)\theta(C_{a35}) + \pi^2 \left(-N_0(\sqrt{|C_{a35}|}|\mathbf{b_2} + \mathbf{b_1'}|) + iJ_0(\sqrt{|C_{a35}|}|\mathbf{b_2} + \mathbf{b_1'}|)\right) \theta(-C_{a35})\right] \\
&[2\pi K_0(\sqrt{D_{a35}}| - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|)\theta(D_{a35}) + \\
&\pi^2 \left(-N_0(\sqrt{|D_{a35}|}| - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|) + iJ_0(\sqrt{|D_{a35}|}| - \mathbf{b_1'} + \mathbf{b_3'} - \mathbf{b_q}|)\right) \theta(-D_{a35})]
\end{aligned}
\tag{210}
$$

$$
\begin{aligned}
&\Omega^{a36} = \\
&\left[2\pi K_0(\sqrt{A_{a36}}|\mathbf{b_2}-\mathbf{b_q}|)\theta(A_{a36}) + \pi^2\left(-N_0(\sqrt{|A_{a36}|}|\mathbf{b_2}-\mathbf{b_q}|) + iJ_0(\sqrt{|A_{a36}|}|\mathbf{b_2}-\mathbf{b_q}|)\right)\theta(-A_{a36})\right] \\
&\left[2\pi K_0(\sqrt{B_{a36}}|\mathbf{b'_2}|)\theta(B_{a36}) + \pi^2\left(-N_0(\sqrt{|B_{a36}|}|\mathbf{b'_2}|) + iJ_0(\sqrt{|B_{a36}|}|\mathbf{b'_2}|)\right)\theta(-B_{a36})\right] \\
&[2\pi K_0(\sqrt{C_{a36}}|\mathbf{b_2}-\mathbf{b'_2}+\mathbf{b'_3}-\mathbf{b_q}|)\theta(C_{a36}) + \\
&\pi^2\left(-N_0(\sqrt{|C_{a36}|}|\mathbf{b_2}-\mathbf{b'_2}+\mathbf{b'_3}-\mathbf{b_q}|) + iJ_0(\sqrt{|C_{a36}|}|\mathbf{b_2}-\mathbf{b'_2}+\mathbf{b'_3}-\mathbf{b_q}|)\right)\theta(-C_{a36})] \\
&\left[2\pi K_0(\sqrt{D_{a36}}|\mathbf{b'_3}-\mathbf{b_q}|)\theta(D_{a36}) + \pi^2\left(-N_0(\sqrt{|D_{a36}|}|\mathbf{b'_3}-\mathbf{b_q}|) + iJ_0(\sqrt{|D_{a36}|}|\mathbf{b'_3}-\mathbf{b_q}|)\right)\theta(-D_{a36})\right]
\end{aligned}
\tag{211}
$$

$$
\begin{aligned}
&\Omega^{b1} = \\
&\int_0^1 \frac{dz_1 dz_2 dz_3}{z_1(1-z_1)z_2(1-z_2)}\frac{\sqrt{X_3}}{\sqrt{|Z_3|}}\left[\pi^3 K_1(\sqrt{X_3 Z_3})\theta(Z_3) + \frac{\pi^4}{2}[N_1(\sqrt{X_3|Z_3|}) - iJ_1(\sqrt{X_3|Z_3|})]\theta(-Z_3)\right]
\end{aligned}
\tag{212}
$$

with

$$
\begin{aligned}
Z_3 &= B_{b1}(1-z_3) + \frac{z_3}{z_2(1-z_2)}\left[C_{b1}(1-z_2) + \frac{z_2}{z_1(1-z_1)}[A_{b1}(1-z_1) + D_{b1}z_1]\right] \\
X_3 &= [-\mathbf{b'_1} + z_2(\mathbf{b'_1}+\mathbf{b_q}) - (\mathbf{b_2}-\mathbf{b_q})z_2(1-z_1)]^2 + \frac{z_2(1-z_2)}{z_3}[-\mathbf{b'_1}-\mathbf{b_q}-(\mathbf{b_2}-\mathbf{b_q})z_1]^2.
\end{aligned}
\tag{213}
$$

$$
\begin{aligned}
&\Omega^{b2} = \\
&\int_0^1 dz\frac{|\mathbf{b'_1}+\mathbf{b_q}|}{\sqrt{|Z_1|}}\left[\pi K_1(\sqrt{Z_1}|\mathbf{b'_1}+\mathbf{b_q}|)\theta(Z_1) + \frac{\pi^2}{2}[N_1(\sqrt{|Z_1|}|\mathbf{b'_1}+\mathbf{b_q}|) - iJ_1(\sqrt{|Z_1|}|\mathbf{b'_1}+\mathbf{b_q}|)]\theta(-Z_1)\right] \\
&\left[2\pi K_0(\sqrt{A_{b2}}|\mathbf{b_2}+\mathbf{b'_1}|)\theta(A_{b2}) + \pi^2\left(-N_0(\sqrt{|A_{b2}|}|\mathbf{b_2}+\mathbf{b'_1}|) + iJ_0(\sqrt{|A_{b2}|}|\mathbf{b_2}+\mathbf{b'_1}|)\right)\theta(-A_{b2})\right] \\
&\left[2\pi K_0(\sqrt{B_{b2}}|\mathbf{b_q}|)\theta(B_{b2}) + \pi^2\left(-N_0(\sqrt{|B_{b2}|}|\mathbf{b_q}|) + iJ_0(\sqrt{|B_{b2}|}|\mathbf{b_q}|)\right)\theta(-B_{b2})\right]
\end{aligned}
\tag{214}
$$

with

$$
Z_1 = C_{b2}z + D_{b2}(1-z). \tag{215}
$$

$$
\begin{aligned}
&\Omega^{b3} = \\
&\left[2\pi K_0(\sqrt{A_{b3}}|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|)\theta(A_{b3}) + \pi^2\left(-N_0(\sqrt{|A_{b3}|}|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|) + iJ_0(\sqrt{|A_{b3}|}|\mathbf{b_2}-\mathbf{b_3}-\mathbf{b_q}|)\right)\theta(-A_{b3})\right] \\
&\left[2\pi K_0(\sqrt{B_{b3}}|\mathbf{b_3}-\mathbf{b'_1}|)\theta(B_{b3}) + \pi^2\left(-N_0(\sqrt{|B_{b3}|}|\mathbf{b_3}-\mathbf{b'_1}|) + iJ_0(\sqrt{|B_{b3}|}|\mathbf{b_3}-\mathbf{b'_1}|)\right)\theta(-B_{b3})\right] \\
&\left[2\pi K_0(\sqrt{C_{b3}}|\mathbf{b'_1}+\mathbf{b_q}|)\theta(C_{b3}) + \pi^2\left(-N_0(\sqrt{|C_{b3}|}|\mathbf{b'_1}+\mathbf{b_q}|) + iJ_0(\sqrt{|C_{b3}|}|\mathbf{b'_1}+\mathbf{b_q}|)\right)\theta(-C_{b3})\right] \\
&\left[2\pi K_0(\sqrt{D_{b3}}|\mathbf{b_3}|)\theta(D_{b3}) + \pi^2\left(-N_0(\sqrt{|D_{b3}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{b3}|}|\mathbf{b_3}|)\right)\theta(-D_{b3})\right]
\end{aligned}
\tag{216}
$$

$$
\begin{aligned}
&\Omega^{b4} = \\
&\Big[2\pi K_0(\sqrt{A_{b4}}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b'_1}|)\theta(A_{b4})+\pi^2\left(-N_0(\sqrt{|A_{b4}|}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b'_1}|)+iJ_0(\sqrt{|A_{b4}|}|\mathbf{b_2}-\mathbf{b_3}+\mathbf{b'_1}|)\right)\theta(-A_{b4})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{b4}}|\mathbf{b_3}+\mathbf{b_q}|)\theta(B_{b4})+\pi^2\left(-N_0(\sqrt{|B_{b4}|}|\mathbf{b_3}+\mathbf{b_q}|)+iJ_0(\sqrt{|B_{b4}|}|\mathbf{b_3}+\mathbf{b_q}|)\right)\theta(-B_{b4})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{b4}}|\mathbf{b'_1}+\mathbf{b_q}|)\theta(C_{b4})+\pi^2\left(-N_0(\sqrt{|C_{b4}|}|\mathbf{b'_1}+\mathbf{b_q}|)+iJ_0(\sqrt{|C_{b4}|}|\mathbf{b'_1}+\mathbf{b_q}|)\right)\theta(-C_{b4})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{b4}}|\mathbf{b_3}|)\theta(D_{b4})+\pi^2\left(-N_0(\sqrt{|D_{b4}|}|\mathbf{b_3}|)+iJ_0(\sqrt{|D_{b4}|}|\mathbf{b_3}|)\right)\theta(-D_{b4})\Big]
\end{aligned}
\tag{217}
$$

$$
\begin{aligned}
&\Omega^{b5} = \\
&\Big[2\pi K_0(\sqrt{A_{b5}}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|)\theta(A_{b5})+\pi^2\left(-N_0(\sqrt{|A_{b5}|}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|)+iJ_0(\sqrt{|A_{b5}|}|\mathbf{b_1}-\mathbf{b_2}+\mathbf{b_q}|)\right)\theta(-A_{b5})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{b5}}|\mathbf{b_1}-\mathbf{b'_1}|)\theta(B_{b5})+\pi^2\left(-N_0(\sqrt{|B_{b5}|}|\mathbf{b_1}-\mathbf{b'_1}|)+iJ_0(\sqrt{|B_{b5}|}|\mathbf{b_1}-\mathbf{b'_1}|)\right)\theta(-B_{b5})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{b5}}|\mathbf{b'_1}+\mathbf{b_q}|)\theta(C_{b5})+\pi^2\left(-N_0(\sqrt{|C_{b5}|}|\mathbf{b'_1}+\mathbf{b_q}|)+iJ_0(\sqrt{|C_{b5}|}|\mathbf{b'_1}+\mathbf{b_q}|)\right)\theta(-C_{b5})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{b5}}|\mathbf{b_1}|)\theta(D_{b5})+\pi^2\left(-N_0(\sqrt{|D_{b5}|}|\mathbf{b_1}|)+iJ_0(\sqrt{|D_{b5}|}|\mathbf{b_1}|)\right)\theta(-D_{b5})\Big]
\end{aligned}
\tag{218}
$$

$$
\begin{aligned}
&\Omega^{b6} = \\
&\Big[2\pi K_0(\sqrt{A_{b6}}|\mathbf{b'_2}|)\theta(A_{b6})+\pi^2\left(-N_0(\sqrt{|A_{b6}|}|\mathbf{b'_2}|)+iJ_0(\sqrt{|A_{b6}|}|\mathbf{b'_2}|)\right)\theta(-A_{b6})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{b6}}|\mathbf{b_1}+\mathbf{b_q}|)\theta(B_{b6})+\pi^2\left(-N_0(\sqrt{|B_{b6}|}|\mathbf{b_1}+\mathbf{b_q}|)+iJ_0(\sqrt{|B_{b6}|}|\mathbf{b_1}+\mathbf{b_q}|)\right)\theta(-B_{b6})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{b6}}|\mathbf{b'_1}+\mathbf{b_q}|)\theta(C_{b6})+\pi^2\left(-N_0(\sqrt{|C_{b6}|}|\mathbf{b'_1}+\mathbf{b_q}|)+iJ_0(\sqrt{|C_{b6}|}|\mathbf{b'_1}+\mathbf{b_q}|)\right)\theta(-C_{b6})\Big] \\
&\Big[2\pi K_0(\sqrt{D_{b6}}|\mathbf{b_1}|)\theta(D_{b6})+\pi^2\left(-N_0(\sqrt{|D_{b6}|}|\mathbf{b_1}|)+iJ_0(\sqrt{|D_{b6}|}|\mathbf{b_1}|)\right)\theta(-D_{b6})\Big]
\end{aligned}
\tag{219}
$$

$$
\begin{aligned}
&\Omega^{b7} = \\
&\Big[2\pi K_0(\sqrt{A_{b7}}|\mathbf{b_2}-\mathbf{b_q}|)\theta(A_{b7})+\pi^2\left(-N_0(\sqrt{|A_{b7}|}|\mathbf{b_2}-\mathbf{b_q}|)+iJ_0(\sqrt{|A_{b7}|}|\mathbf{b_2}-\mathbf{b_q}|)\right)\theta(-A_{b7})\Big] \\
&\Big[2\pi K_0(\sqrt{B_{b7}}|\mathbf{b'_1}|)\theta(B_{b7})+\pi^2\left(-N_0(\sqrt{|B_{b7}|}|\mathbf{b'_1}|)+iJ_0(\sqrt{|B_{b7}|}|\mathbf{b'_1}|)\right)\theta(-B_{b7})\Big] \\
&\Big[2\pi K_0(\sqrt{C_{b7}}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|)\theta(C_{b7})+\pi^2\left(-N_0(\sqrt{|C_{b7}|}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|)+iJ_0(\sqrt{|C_{b7}|}|\mathbf{b_2}+\mathbf{b'_1}-\mathbf{b'_2}|)\right)\theta(-C_{b7})\Big] \\
&[2\pi K_0(\sqrt{D_{b7}}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|)\theta(D_{b7})+ \\
&\pi^2\left(-N_0(\sqrt{|D_{b7}|}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|)+iJ_0(\sqrt{|D_{b7}|}|-\mathbf{b_2}+\mathbf{b'_2}+\mathbf{b_q}|)\right)\theta(-D_{b7})]
\end{aligned}
\tag{220}
$$

$$
\begin{aligned}
&\Omega^{b8} = \\
&\left[2\pi K_0(\sqrt{A_{b8}}|\mathbf{b_2'}|)\theta(A_{b8}) + \pi^2\left(-N_0(\sqrt{|A_{b8}|}|\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{b8}|}|\mathbf{b_2'}|)\right)\theta(-A_{b8})\right] \\
&[2\pi K_0(\sqrt{B_{b8}}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|)\theta(B_{b8}) + \\
&\pi^2\left(-N_0(\sqrt{|B_{b8}|}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{b8}|}|-\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}+\mathbf{b_q}|)\right)\theta(-B_{b8})] \\
&\left[2\pi K_0(\sqrt{C_{b8}}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|)\theta(C_{b8}) + \pi^2\left(-N_0(\sqrt{|C_{b8}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|) + iJ_0(\sqrt{|C_{b8}|}|\mathbf{b_2}+\mathbf{b_1'}-\mathbf{b_2'}|)\right)\theta(-C_{b8})\right] \\
&[2\pi K_0(\sqrt{D_{b8}}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|)\theta(D_{b8}) + \\
&\pi^2\left(-N_0(\sqrt{|D_{b8}|}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|) + iJ_0(\sqrt{|D_{b8}|}|-\mathbf{b_2}+\mathbf{b_2'}+\mathbf{b_q}|)\right)\theta(-D_{b8})]
\end{aligned}
\tag{221}
$$

$$
\begin{aligned}
&\Omega^{c1} = \\
&\frac{1}{2}\left[2\pi K_0(\sqrt{A_{c1}}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}|)\theta(A_{c1}) + \pi^2\left(-N_0(\sqrt{|A_{c1}|}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}|) + iJ_0(\sqrt{|A_{c1}|}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}+\mathbf{b_2'}|)\right)\theta(-A_{c1})\right] \\
&\left[2\pi K_0(\sqrt{B_{c1}}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_2'}|)\theta(B_{c1}) + \pi^2\left(-N_0(\sqrt{|B_{c1}|}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_2'}|) + iJ_0(\sqrt{|B_{c1}|}\frac{1}{2}|\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_2'}|)\right)\theta(-B_{c1})\right] \\
&[2\pi K_0(\sqrt{C_{c1}}\frac{1}{2}|-\mathbf{b_2}+\mathbf{b_1'}+\mathbf{b_2'}+2\mathbf{b_q}|)\theta(C_{c1}) + \\
&\pi^2\left(-N_0(\sqrt{|C_{c1}|}\frac{1}{2}|-\mathbf{b_2}+\mathbf{b_1'}+\mathbf{b_2'}+2\mathbf{b_q}|) + iJ_0(\sqrt{|C_{c1}|}\frac{1}{2}|-\mathbf{b_2}+\mathbf{b_1'}+\mathbf{b_2'}+2\mathbf{b_q}|)\right)\theta(-C_{c1})] \\
&\left[2\pi K_0(\sqrt{D_{c1}}|\mathbf{b_2}-\mathbf{b_2'}-\mathbf{b_q}|)\theta(D_{c1}) + \pi^2\left(-N_0(\sqrt{|D_{c1}|}|\mathbf{b_2}-\mathbf{b_2'}-\mathbf{b_q}|) + iJ_0(\sqrt{|D_{c1}|}|\mathbf{b_2}-\mathbf{b_2'}-\mathbf{b_q}|)\right)\theta(-D_{c1})\right]
\end{aligned}
\tag{222}
$$

$$
\begin{aligned}
&\Omega^{c2} = \\
&\frac{1}{2}[2\pi K_0(\sqrt{A_{c2}}\frac{1}{2}|2\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_q}|)\theta(A_{c2}) + \\
&\pi^2\left(-N_0(\sqrt{|A_{c2}|}\frac{1}{2}|2\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_q}|) + iJ_0(\sqrt{|A_{c2}|}\frac{1}{2}|2\mathbf{b_2}-\mathbf{b_1'}-\mathbf{b_q}|)\right)\theta(-A_{c2})] \\
&[2\pi K_0(\sqrt{B_{c2}}\frac{1}{2}|2\mathbf{b_3}-\mathbf{b_1'}+\mathbf{b_q}|)\theta(B_{c2}) + \\
&\pi^2\left(-N_0(\sqrt{|B_{c2}|}\frac{1}{2}|2\mathbf{b_3}-\mathbf{b_1'}+\mathbf{b_q}|) + iJ_0(\sqrt{|B_{c2}|}\frac{1}{2}|2\mathbf{b_3}-\mathbf{b_1'}+\mathbf{b_q}|)\right)\theta(-B_{c2})] \\
&\left[2\pi K_0(\sqrt{C_{c2}}\frac{1}{2}|\mathbf{b_1'}+2\mathbf{b_q}|)\theta(C_{c2}) + \pi^2\left(-N_0(\sqrt{|C_{c2}|}\frac{1}{2}|\mathbf{b_1'}+2\mathbf{b_q}|) + iJ_0(\sqrt{|C_{c2}|}\frac{1}{2}|\mathbf{b_1'}+2\mathbf{b_q}|)\right)\theta(-C_{c2})\right] \\
&\left[2\pi K_0(\sqrt{D_{c2}}|\mathbf{b_3}|)\theta(D_{c2}) + \pi^2\left(-N_0(\sqrt{|D_{c2}|}|\mathbf{b_3}|) + iJ_0(\sqrt{|D_{c2}|}|\mathbf{b_3}|)\right)\theta(-D_{c2})\right]
\end{aligned}
\tag{223}
$$

Appendix E: Expressions for H_F^{ij}

H_F^{ij} corresponding to the form factors defined in eq.(2) with a "tilde" for $\overline{B^0} \to \Lambda_c^+ \overline{p}$. The expressions for diagrams (a1)∼(a36) in Fig. 1, (b1)∼(b8) in Fig. 2 and (c1), (c2) in Fig. 3 are listed in the following.

For the hard amplitudes of Fig.1(a1):

$$
\begin{aligned}
&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^5 r^2(-1+r^2)(1-y) \\
&\widetilde{A^T} = -\widetilde{B^T} = 32m_B^5 r^2(-1+r^2)(1-y)
\end{aligned}
\tag{224}
$$

For the hard amplitudes of Fig.1(a2):

$$\widetilde{A^V}=-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=16m_B^5r(-1+r^2)^2x_3'(-1+x_2) \tag{225}$$

For the hard amplitudes of Fig.1(a3):

$$\begin{aligned}&\widetilde{A^V}=\widetilde{A^A}=\widetilde{B^V}=\widetilde{B^A}=0\\&\widetilde{A^T}=\widetilde{B^T}=0\end{aligned} \tag{226}$$

For the hard amplitudes of Fig.1(a4):

$$\begin{aligned}&\widetilde{A^V}=\widetilde{A^A}=\widetilde{B^V}=\widetilde{B^A}=0\\&\widetilde{A^T}=\widetilde{B^T}=0\end{aligned} \tag{227}$$

For the hard amplitudes of Fig.1(a5):

$$\widetilde{A^V}=-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=-16m_B^5r(r^2-1)^2x_2x_3' \tag{228}$$

For the hard amplitudes of Fig.1(a6):

$$\begin{aligned}&\widetilde{A^V}=\widetilde{B^A}=-16m_B^5r(-1+r^2)(r(-1+x_2+y)+(-1+x_2)(x_2'-r^2x_2'+r(-1+x_2+y)))\\&\widetilde{A^A}=\widetilde{B^V}=-16m_B^5r(-1+r^2)(r(-1+x_2+y)+(-1+x_2)(-x_2'+r^2x_2'+r(-1+x_2+y)))\\&\widetilde{A^T}=-\widetilde{B^T}=-32m_B^5r^2(-1+r^2)x_2(-1+x_2+y)\end{aligned} \tag{229}$$

For the hard amplitudes of Fig.1(a7):

$$\widetilde{A^V}=-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=-16m_B^5r(-1+r^2)x_2 \tag{230}$$

For the hard amplitudes of Fig.1(a8):

$$\begin{aligned}\widetilde{A^V}\;&=\;\widetilde{B^A}=-16m_B^4(1-r^2)(m_cr+m_B(r^2(-1+x_3')-x_3')+(x_2-y)(-(m_c(-1+r)r)+\\&\quad m_B(r^2(-1+x_3')-x_3'+r^3(-1+x_1+y))))\\\widetilde{A^A}\;&=\;\widetilde{B^V}=16m_B^4(1-r^2)(-m_cr+m_B(-r^2(-1+x_3')+x_3')+(x_2-y)(-(m_cr(1+r))+\\&\quad m_B(-(r^2(-1+x_3'))+x_3'+r^3(-1+x_1+y))))\end{aligned} \tag{231}$$

For the hard amplitudes of Fig.1(a9):

$$\begin{aligned}\widetilde{A^V}\;&=\;-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=16m_B^3(-1+r^2)(m_c^2r+m_cm_B(r^2(-1+x_3')-x_3')+\\&\quad m_cm_B(1+(-1+r^2)x_1'+r^2(-x_3+y))-m_B^2r(-1+x_1+y)(1+(-1+r^2)x_1'+r^2(-x_3+y)))\end{aligned} \tag{232}$$

For the hard amplitudes of Fig.1(a10):

$$\widetilde{A^V}\;=\;-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=16m_cm_B^4r^2(-1+r^2)(1-y) \tag{233}$$

For the hard amplitudes of Fig.1(a11):

$$\widetilde{A^V}\;=\;-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=-16m_B^4(-1+r^2)^2x_2'(-m_c+m_Br(-1+x_1+y)) \tag{234}$$

For the hard amplitudes of Fig.1(a12):

$$\widetilde{A^V}\;=\;\widetilde{A^A}=\widetilde{B^V}=\widetilde{B^A}=-16m_B^4(-1+r^2)x_2(m_cr+m_B(r^2(-1+x_3')-x_3')) \tag{235}$$

For the hard amplitudes of Fig.1(a13):

$$\widetilde{A^V}\;=\;-\widetilde{A^A}=-\widetilde{B^V}=\widetilde{B^A}=-16m_B^4(1-r^2)(m_Br^3(x_3-y)(-1+x_2+y)+m_c(x_2'+r^2(-1+x_2-x_2'+y))) \tag{236}$$

For the hard amplitudes of Fig.1(a14):

$$\widetilde{A^V} \;=\; -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5 r(-1+r^2)(1+(-1+r^2)x_1')x_2 \tag{237}$$

For the hard amplitudes of Fig.1(a15):

$$\begin{aligned}\widetilde{A^V} &= \widetilde{B^A} = 16m_B^5(-1+r^2)((1-r^2)x_3' - (x_2-y)(-x_3'+r^2x_3'+r^3(-1+x_3+y)))\\ \widetilde{A^A} &= \widetilde{B^V} = -16m_B^5(-1+r^2)((-1+r^2)x_3' - (x_2-y)(x_3'-r^2x_3'+r^3(-1+x_3+y)))\end{aligned} \tag{238}$$

For the hard amplitudes of Fig.1(a16):

$$\widetilde{A^V} \;=\; -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_c m_B^4 r^2(-1+r^2)(-1+x_3+y) \tag{239}$$

For the hard amplitudes of Fig.1(a17):

$$\widetilde{A^V} \;=\; -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^5 r^3(-1+r^2)(-x_1+y)(-1+x_3+y) \tag{240}$$

For the hard amplitudes of Fig.1(a18):

$$\widetilde{A^V} \;=\; -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5 r(-1+r^2)^2 x_2'(x_1-y) \tag{241}$$

For the hard amplitudes of Fig.1(a19):

$$\widetilde{A^V} \;=\; \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^5(-1+r^2)^2 x_2 x_3' \tag{242}$$

For the hard amplitudes of Fig.1(a20):

$$\widetilde{A^V} \;=\; -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5 r(1-r^2)(-x_1+y)(x_2'+r^2(-1+x_2-x_2'+y)) \tag{243}$$

For the hard amplitudes of Fig.1(a21):

$$\begin{aligned}&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 0\\ &\widetilde{A^T} = \widetilde{B^T} = 0\end{aligned} \tag{244}$$

For the hard amplitudes of Fig.1(a22):

$$\begin{aligned}&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = -16m_B^5 r^2(-1+r^2)(1-y)\\ &\widetilde{A^T} = -\widetilde{B^T} = -32m_B^5 r^2(-1+r^2)(1-y)\end{aligned} \tag{245}$$

For the hard amplitudes of Fig.1(a23):

$$\begin{aligned}&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^4(-1+r^2)(m_c r + m_B(-x_2'+r^2(-2+x_2'+x_3)))(1-y)\\ &\widetilde{A^T} = -\widetilde{B^T} = -32m_B^4 r(-1+r^2)(2m_c+m_B r(-1+x_3))(1-y)\end{aligned} \tag{246}$$

For the hard amplitudes of Fig.1(a24):

$$\begin{aligned}&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = -16m_B^5(-1+r^2)^2 x_2'(1-y)\\ &\widetilde{A^T} = -\widetilde{B^T} = 64m_B^5 r^2(-1+r^2)(-1+x_1)(1-y)\end{aligned} \tag{247}$$

For the hard amplitudes of Fig.1(a25):

$$\begin{aligned}&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^5 r^2(-1+r^2)(1-y)(-1+x_2+y)\\ &\widetilde{A^T} = -\widetilde{B^T} = 32m_B^5 r^2(-1+r^2)(1-y)(-1+x_2+y)\end{aligned} \tag{248}$$

For the hard amplitudes of Fig.1(a26):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^5 r(-1+r^2)^2 x_2 \tag{249}$$

For the hard amplitudes of Fig.1(a27):

$$\begin{aligned}
&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^5r^2(-1+r^2)(-1+x_2+y)^2\\
&\widetilde{A^T} = -\widetilde{B^T} = 32m_B^5r^2(-1+r^2)(-1+x_2+y)^2
\end{aligned} \tag{250}$$

For the hard amplitudes of Fig.1(a28):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5r(-1+r^2)^2x_2 \tag{251}$$

For the hard amplitudes of Fig.1(a29):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^4r(-1+r^2)(1-y)(-m_cr+m_B(1+(-1+r^2)x_1'+r^2(-x_3+y))) \tag{252}$$

For the hard amplitudes of Fig.1(a30):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5r^3(-1+r^2)(1-y)(-x_1+y) \tag{253}$$

For the hard amplitudes of Fig.1(a31):

$$\begin{aligned}
&\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^5r(-1+r^2)(1-y)\\
&\widetilde{A^T} = -\widetilde{B^T} = 32m_B^5r^2(-1+r^2)(1-y)x_2
\end{aligned} \tag{254}$$

For the hard amplitudes of Fig.1(a32):

$$\begin{aligned}
&\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^5r(-1+r^2)^2(1-y)\\
&\widetilde{A^T} = -\widetilde{B^T} = -32m_B^5r^2(-1+r^2)(1-y)(-1+x_2+y)
\end{aligned} \tag{255}$$

For the hard amplitudes of Fig.1(a33):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^4m_c(-1+r^2)^2x_3' \tag{256}$$

For the hard amplitudes of Fig.1(a34):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5r(-1+r^2)^2x_3'(-x_1+y) \tag{257}$$

For the hard amplitudes of Fig.1(a35):

$$\begin{aligned}
&\widetilde{A^V} = \widetilde{B^A} = -16m_B^5r(1-r^2)(r(-1+x_2+y)+(-1+x_2)(-x_3'+r^2x_3'+r(-1+x_2+y)))\\
&\widetilde{A^A} = \widetilde{B^V} = -16m_B^5r(1-r^2)(r(-1+x2+y)+(-1+x_2)(x_3'-r^2x_3'+r(-1+x_2+y)))
\end{aligned} \tag{258}$$

For the hard amplitudes of Fig.1(a36):

$$\widetilde{A^V} = \widetilde{B^A} = \widetilde{A^A} = \widetilde{B^V} = -16m_B^5r^2(-1+r^2)(-1+x_2+y)^2 \tag{259}$$

For the hard amplitudes of Fig.2(b1):

$$\begin{aligned}
&\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 0\\
&\widetilde{A^T} = \widetilde{B^T} = 0
\end{aligned} \tag{260}$$

For the hard amplitudes of Fig.2(b2):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -16m_B^5r(-1+r^2)x_2^2 \tag{261}$$

For the hard amplitudes of Fig.2(b3):

$$\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = -16m_B^4(-1+r^2)x_2(-m_cr+m_B((-1+r^2)x_1'+y)) \tag{262}$$

For the hard amplitudes of Fig.2(b4):

$$\widetilde{A^T} = -\widetilde{B^T} = 32m_B^4 r(-1+r^2)x_2(m_c + m_B r(-1+x_3)) \tag{263}$$

For the hard amplitudes of Fig.2(b5):

$$\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 16m_B^5(-1+r^2)x_2(r^2(-1+x_1') - x_1' + y) \tag{264}$$

For the hard amplitudes of Fig.2(b6):

$$\widetilde{A^T} = -\widetilde{B^T} = -32m_B^5 r^2(-1+r^2)x_2(-1+x_1) \tag{265}$$

For the hard amplitudes of Fig.2(b7):

$$\begin{aligned} &\widetilde{A^V} = \widetilde{A^A} = \widetilde{B^V} = \widetilde{B^A} = 0 \\ &\widetilde{A^T} = \widetilde{B^T} = 0 \end{aligned} \tag{266}$$

For the hard amplitudes of Fig.2(b8):

$$\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = 16m_B^5 r(-1+r^2)x_2^2 \tag{267}$$

For the hard amplitudes of Fig.3(c1):

$$\begin{aligned} \widetilde{A^V} &= \widetilde{B^A} = 8m_B^5 r(1-r^2)(1+x_2-y-r(-2+x_2+2y)-(-1+x_2)(x_2'-x_3'+r^2(-x_2'+x_3')+r(-2+x_2+2y))) \\ \widetilde{A^A} &= \widetilde{B^V} = -8m_B^5 r(1-r^2)(1+x_2-y+r(-2+x_2+2y)+(-1+x_2)(-x_2'+r^2(x_2'-x_3')+x_3'+r(-2+x_2+2y))) \\ \widetilde{A^T} &= -\widetilde{B^T} = 16m_B^5 r^2(-1+r^2)x_2(-2+x_2+2y) \end{aligned} \tag{268}$$

For the hard amplitudes of Fig.3(c2):

$$\begin{aligned} &\widetilde{A^V} = -\widetilde{A^A} = -\widetilde{B^V} = \widetilde{B^A} = -8m_B^4(-1+r^2)(m_B r((1+(-1+r^2)x_1'+r^2(x_3-y))(1+x_2-y)+ \\ &3r^2(x_3-y)(-1+y))+m_c(x_2'-x_3'+r^2(-2+x_2-x_2'+x_3'+2y))) \end{aligned} \tag{269}$$

[1] ARGUS Collaboration, H. Albrecht, Phys. Lett. **B209** (1988) 119.

[2] Babar Collaborqtion, B. Aubert et al., Phys. Rev. **D69**, 091503(2004); Phys. **D72**, 051101(2005); Phys. Rev. Lett. **90**, 231801(2003); Belle Collaboration, M.C. Chang et al., Phys. Rev. **D71**, 072007(2005); Q.L. Xie et al., Phys. Rev. **D72**, 051105(2005); M.Z. Wang et al., Phys. Rev. Lett. **90**, 201802(2003); ibid. **92**, 131801(2004); Y.J. Lee et al., Phys. Rev. Lett. **93**, 211801(2004); M.Z. Wang et al., Phys. Lett. **B617**, 141(2995); K. Abe et al., Phys. Rev. Lett. **88**, 181803(2005); CLEO Collaborqtion, A. Bornheim et al., Phys. Rev. **D68**, 052002(2003); T. E. Coan et al., ibid **59**, 111101(1999); S. Anderson et al., Phys. Rev. Lett. **86**, 2732(2001).

[3] Belle Collaboration, N. Gabyshev et al., hep-ex/0409005; Belle Collaboration, N. Gabyshev et al., Phys. Rev. Lett. **90** (2003) 121802; Belle Collaboration, R. Chistov et al., hep-ex/0510074.

[4] Babar Collaboration, B. Aubert et al., hep-ex/0607055; T. Berger-Hryn'ova, "Study of B meson decays to p $\bar{p}$ h final states", Ph.D. thesis, SLAC-R-810.

[5] CLEO Collaboration, S.A. Dytman et al., Phys. Rev. **D66** (2002) 091101.

[6] S. Eidelman, et al., Particle Data Group, Phys. Lett. **B592** (2004) 1.

[7] Yong-Yeon Keum and Hsiang-nan Li, Phys. Rev. **D63** (2001) 074006; Chuan-Hung Chen, Yong-Yeon Keum and Hsiang-nan Li, Phys. Rev. **D64** (2001) 112002; Yong-Yeon Keum, Hsiang-nan Li and A.I. Sanda, Phys. Rev. **D63** (2001) 054008; Chuan-Hung Chen, Yong-Yeon Keum and Hsiang-nan Li, Phys. Rev. **D66** (2002) 054013; Cai-Dian Lü, Kazumasa Ukai and Mao-Zhi Yang, Phys. Rev. **D63** (2001) 074009; Cai-Dian Lü and Mao-Zhi Yang, Eur. Phys. J. **C23** (2002) 275.

[8] Xia-Gang He, B. McKellar and D.-D. Wu, Phys. Rev. **D41**, 2141(1990).

[9] V. Chernyak and I. Zhitnitsky, Nuch. Phys. **B345** (1990) 137.

[10] G. Lu, X. Xue and J. Liu, Phys. Lett. **B259** (1991) 169.

[11] M. Jarfi et al., Phys. Rev. **D43** (1991) 1599; Phys. Lett. **B237** (1990) 513.

[12] P. Ball and H.G. Dosch, Z. Phys. **C51** (1991) 169.

[13] H.Y. Cheng and K.C. Yang, Phys. Rev. **D67** (2003) 034008; ibid. **65** (2002) 054028.

[14] H.Y. Cheng, talk presented at 3rd International Conference on Flavor Physics, Oct 3-8, 2005, National Central Univ. Chung-li, Taiwan, hep-ph/0603003.

[15] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125.

[16] Hsien-Hung Shih, Shih-Chang Lee and Hsiang-nan Li, Phys. Rev. **D59** (1999) 094014.

[17] Hsien-Hung Shih, Shih-Chang Lee and Hsiang-nan Li, Phys. Rev. **D61** (2000) 114002.

[18] Chung-Hsien Chou, Hsien-Hung Shih, Shih-Chang Lee and Hsiang-nan Li, Phys. Rev. **D65** (2002) 074030.

[19] W. Loinaz and R. Akhoury, Phys. Rev. **D53** (1996) 1416; F. Schlumpf, Ph.D. Thesis, Zurich University, 1992, arXiv:hep-ph/9211255.

[20] V.L. Chernyak, A.A. Ogloblin and I.R. Zhitnitsky, Z. Phys. **C42** (1989) 569.

[21] A.G. Grozin and M. Neubert, Phys. Rev. **D55** (1997) 272; M. Beneke and T. Feldmann, Nucl. Phys. **B592** (2001) 3; M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Nucl. Phys. **B591** (2000) 313; T. Kurimoto, Hsiang-nan Li and A. I. Sanda, Phys. Rev. **D65** (2001) 014007.

[22] DELPHI Collaboration, J. Abdallah et al., Phys. Lett. **B585** (2004) 63.

[23] Xiao-Gang He, Tong Li, Xue-Qian Li and Yu-Ming Wang, Phys. Rev. **D74** (2006) 034026.

[24] Yong-Yeon Keum, T. Kurimoto, Hsiang-nan Li, Cai-Dian Lü and A. I. Sanda, Phys. Rev. **D69** (2004) 094018.

[25] Bijoy Kundu, Hsiang-nan Li, Jim Samuelsson and Pankaj Jain, Eur. Phys. J. **C8** (1999) 637.